

EQUIVARIANT ELLIPTIC GENERA AND LOCAL MCKAY CORRESPONDENCES

ROBERT WAELDER

ABSTRACT. In this paper we prove an equivariant version of the McKay correspondence for the elliptic genus on open varieties with a torus action. As a consequence, we will prove the equivariant DMVV formula for the Hilbert scheme of points on $\mathbb{C}^2$.

1. INTRODUCTION

Let X be a smooth variety, and G a finite group acting holomorphically on X . Suppose that the quotient X/G possesses a crepant resolution of singularities V . The McKay correspondence refers to the identification of topological invariants of V with orbifold analogues of these invariants associated to the action of G on X . The classical example is the case in which $G \subset SU(2)$ is a finite subgroup acting on $\mathbb{C}^2$. Then the quotient $\mathbb{C}^2/G$ possesses a unique crepant resolution V , and the Euler characteristic of V coincides with the orbifold Euler characteristic $e_{orb}(\mathbb{C}^2, G) = \frac{1}{|G|} \sum_{gh=hg} e((\mathbb{C}^2)^{g,h})$. Over the years, the McKay correspondence has exhibited a remarkable versatility towards generalization. In [5] Batyrev investigated a more general class of resolutions $\widehat{X}/G \rightarrow X/G$, and proved the McKay correspondence for the E -function in this situation. More recently, Borisov and Libgober have proven a similar result for the elliptic genus [7].

In this paper we will prove an equivariant analogue of the McKay Correspondence for the elliptic genus. The advantage of working in the equivariant setting is that, by localization, we can make sense of the elliptic genus even for open varieties. This allows us to prove a host of new formulas. One consequence of the work in this paper is a beautiful formula for the generating function of the equivariant elliptic genus of the Hilbert scheme of points on $\mathbb{C}^2$ (with the standard torus action):

$$\sum_{n \geq 0} p^n Ell((\mathbb{C}^2)^{[n]}; y, q, t_1, t_2) = \prod_{m \geq 0, n \geq 0, \ell, k} \frac{1}{(1 - p^n q^m y^\ell t_1^{k_1} t_2^{k_2})^{c(nm, \ell, k)}}.$$

The terms $c(m, \ell, k)$ are the coefficients in the expansion of $Ell(\mathbb{C}^2; y, q, t_1, t_2)$ in y, q, t_1 , and t_2 . The above formula is an equivariant generalization of the

DMVV formula:

$$\sum_{n>0} p^n \text{Ell}(S^{[n]}; y, q) = \prod_{m \geq 0, n > 0, \ell, k} \frac{1}{(1 - p^n q^m y^\ell)^{c(nm, \ell, k)}}.$$

In the above formula, S is a compact algebraic surface, and $\text{Ell}(S^{[n]}; y, q)$ is the elliptic genus of the Hilbert scheme of n points on S . The non-equivariant DMVV formula was conjectured by string theorists Dijkgraaf, Moore, Verlinde and Verlinde [8], and proven by Borisov and Libgober [7]. The equivariant version is a conjecture of Li, Liu, and Zhou [14].

1.1. Background on the Elliptic Genus. For X a smooth complex manifold, the elliptic genus of X is defined as:

$$(1) \quad \text{Ell}(X) = \int_X \prod \frac{x_j \theta(\frac{x_j}{2\pi i} - z, \tau)}{\theta(\frac{x_j}{2\pi i}, \tau)}$$

The product is taken over the Chern roots of the holomorphic tangent bundle to X . $\theta(\cdot, \tau)$ is the Jacobi theta function, and z represents a formal parameter. Setting $y = e^{2\pi i z}$ and $q = e^{2\pi i \tau}$, the elliptic genus may also be interpreted as the index of the following differential operator:

$$(2) \quad y^{-d/2} \bar{\partial} \otimes \bigotimes_{n=1}^{\infty} \Lambda_{-yq^{n-1}} T^* X \otimes \Lambda_{-y^{-1}q^n} T X \otimes S_{q^n} T^* X \otimes S_{q^n} T X$$

The modular properties of the Jacobi theta function endow the elliptic genus with a rich amount of structure. For example, if X is Calabi-Yau, then the elliptic genus is a weak Jacobi form as a function of $(z, \tau) \in \mathbb{C} \times \mathbb{H}$. If X is Calabi-Yau and possesses a nontrivial torus action, Liu has shown that the modular properties of the equivariant index of 2 actually imply its rigidity [15]. In addition to these properties, the elliptic genus encodes a large number of classical algebraic and topological invariants of the space. For example, letting $q \rightarrow 0$ in the expression for the elliptic genus produces $y^{-d/2}$ times the Hirzebruch χ_{-y} genus, whereas letting $y \rightarrow 1$ produces the Euler characteristic of the space.

In [16], [6], Chin-Lung Wang, Borisov, and Libgober investigated the following relative version of the elliptic genus for pairs (X, D) , where $D = \sum_i a_i D_i$ is a smooth divisor with normal crossings and coefficients $a_i \neq 1$:

$$\text{Ell}(X, D) = \int_X \prod \frac{x_j \theta(\frac{x_j}{2\pi i} - z) \frac{\theta'(0)}{2\pi i}}{\theta(\frac{x_j}{2\pi i}) \theta(-z)} \prod_i \frac{\theta(\frac{c_1(D_i)}{2\pi i} - (-a_i + 1)z) \theta(z)}{\theta(\frac{c_1(D_i)}{2\pi i} - z) \theta((-a_i + 1)z)}$$

The modular properties of the Jacobi theta function imply that the relative elliptic genus satisfies the following change of variables formula for blow-up

morphisms: If $f : \tilde{X} \rightarrow X$ is the blow-up of X along a smooth base with normal crossings with respect to the components of D , and $\tilde{D}$ is the divisor on $\tilde{X}$ satisfying: $K_{\tilde{X}} + \tilde{D} = f^*(K_X + D)$, then $Ell(\tilde{X}, \tilde{D}) = Ell(X, D)$. For Z a $\mathbb{Q}$ -Gorenstein variety with log-terminal singularities, and $X \rightarrow Z$ a resolution of singularities with exceptional locus a normal crossing divisor D , Borisov and Libgober define the singular elliptic genus of Z to be the relative elliptic genus of (X, D) . The change of variable formula, together with the Weak Factorization Theorem [1] implies that this definition is well-defined. Moreover, when Z possesses a crepant resolution V , the singular elliptic genus of Z is easily seen to coincide with the elliptic genus of V .

The singular elliptic genus (and its orbifold analogue) plays a crucial role in Borisov and Libgober's proof of the McKay correspondence for elliptic genera. Its utility stems from the fact that it behaves well with respect to a large class of birational modifications. The added flexibility obtained from studying the singular elliptic genus allowed Borisov and Libgober to reduce their proof to calculations involving toroidal embeddings. Their approach is similar in spirit to that of Batyrev in [5], who proved the McKay correspondence for the E -function by using the change of variables formula from motivic integration to reduce the case to calculations on toric varieties.

When X has a nontrivial torus action, we may define the equivariant elliptic genus of X to be the equivariant index of the operator defined in 2. By the index theorem [4], this is the same as the integral in equation 1 obtained by replacing the Chern roots of TX with their equivariant analogues. Similarly, we may define an equivariant version of the relative elliptic genus by replacing appearances of $c_1(D_i)$ with their equivariant extensions.

The bulk of this paper is devoted to proving the change of variables formula for the equivariant orbifold elliptic genus (this case subsumes the non-orbifold case). Once we establish the change of variables formula in this situation, the remaining steps in the proof of the equivariant McKay correspondence for the elliptic genus follow closely the steps given in [7].

1.2. Outline of the Proof. In a recent preprint [17], I proved the equivariant change of variable formula for blow-ups along complete intersections $W = D_1 \cap \dots \cap D_k \subset X$. The idea was to interpret the blow-down $\tilde{X} \rightarrow X$ as a toroidal morphism. The stratification defined by the divisors D_i determined the toroidal structure of X , whereas the stratification defined by the proper transforms of these divisors, together with the exceptional divisor, determined the toroidal structure of $\tilde{X}$. The comparison of the relative elliptic genera of the base space and its blow-up was ultimately reduced to a computation involving the combinatorics of the polyhedral complexes associated to the two toroidal embeddings. This idea was inspired by Borisov and

Libgober’s use of polyhedral complexes in [7] to compute the push-forward of the orbifold elliptic class under the global quotient map.

Later it became apparant that the proof given in [17] could be adapted to the case in which X was a “normal cone space”, i.e., a fiber product of spaces $\mathbb{P}(F \oplus 1)$, where $F \rightarrow W$ was a holomorphic vectorbundle. The idea was that the Chern roots of the tautological quotient bundle $Q_F \rightarrow X$ should play the role of the “divisors” in a polyhedral complex associated to X . Similarly, if $f : \tilde{X} \rightarrow X$ was the blow-up of X along W with exceptional divisor E , then the Chern roots of $f^*Q_F \otimes \mathcal{O}(-E)$, and of $\mathcal{O}(E)$ should behave like the “divisors” of a polyhedral complex associated to $\tilde{X}$. In this paper, we refer to such polyhedral complexes associated to the data of Chern roots as “twisted polyhedral complexes.” The case of a general blow-up may be reduced to cases of this nature by using an equivariant version of deformation to the normal cone.

The breakdown of the sections in this paper are as follows: For generic specializations of the parameters (z, τ) , the integrand of the equivariant elliptic genus is a power series in the equivariant parameters with differential form coefficients. In section 2 we discuss convergence issues related to power series of this type and put their corresponding cohomology theory on solid ground. In section 3 we define our principle objects of study; namely the equivariant orbifold elliptic class and its relative version. In sections 4 and 5 we review some facts about the equivariant cohomology of toric varieties and discuss how it relates to computing equivariant push-forwards of toroidal morphisms. In section 6 we use these results to compute the pushforward of the orbifold elliptic class under a toroidal morphism which is birational to a quotient by a finite group. This result is the equivariant analogue of Lemma 5.4 in [7]. Sections 7 to 10 are devoted to the proof of the equivariant change of variable formula. In section 7, we prove an equivariant analogue of deformation to the normal cone, tailored specifically to handle cohomological data like the orbifold elliptic class. As stated above, this will allow us to reduce the proof of the change of variable formula to the case when X is a normal cone space. In section 8 we prove for completeness a number of technical lemmas regarding spaces of this form. In section 9 we introduce the twisted polyhedral complex for normal cone spaces. In 10 we apply the techniques from the preceding sections to prove the equivariant change of variables formula. Finally, in 11 we prove the equivariant McKay correspondence for elliptic genera, and the equivariant DMVV formula.

Acknowledgements 1. I wish to thank my advisor Professor Kefeng Liu for introducing me to elliptic genera and for his constant support, as well as Professor Anatoly Libgober for his feedback and help with technical aspects of his work.

2. EQUIVARIANT COHOMOLOGY AND POWER SERIES

2.1. Preliminaries on Equivariant Cohomology. We begin by reviewing some basic aspects of equivariant cohomology. For a thorough reference on the subject see [3].

Let M be a smooth manifold and T a torus acting smoothly on M . Let $e_1, \dots, e_\ell$ form a basis for the Lie algebra of T which is dual to the linear forms $u_1, \dots, u_\ell$. Every $X \in \mathfrak{t}$ defines a vectorfield X on M by the formula $X(p) = \frac{d}{dt}|_{t=0} \exp(tX) \cdot p$. Define $\Omega_T^*(M)$ to be the ring of differential forms on M which are annihilated by $\mathcal{L}_X$ for every $X \in \mathfrak{t}$. If we let $d_{\mathfrak{t}} = d + \sum_{\alpha=1}^{\ell} u_{\alpha} i_{e_{\alpha}}$, then $d_{\mathfrak{t}}$ defines an operator on $\Omega_T^*(M) \otimes \mathbb{C}[u_1, \dots, u_{\ell}]$ and satisfies $d_{\mathfrak{t}}^2 = 0$. The Cartan model for equivariant cohomology is defined to be:

$$H_T^*(M)_{\text{Cartan}} = \frac{\ker d_{\mathfrak{t}}}{\text{im } d_{\mathfrak{t}}}.$$

The translation of concepts from cohomology to equivariant cohomology is more or less routine. For example, a T -map $f : M \rightarrow N$ induces a pullback $f^* : H_T^*(Y) \rightarrow H_T^*(X)$ as in ordinary cohomology. Similarly, for any $E \in K_T(X)$ we may define equivariant characteristic classes of E which are equivariant extensions of the ordinary characteristic classes.

If p is a single point with trivial T -action, the equivariant map $\pi : M \rightarrow p$ induces a map $\pi^* : H_T^*(p) \rightarrow H_T^*(M)$. Since $H_T^*(p) = \mathbb{C}[u_1, \dots, u_{\ell}]$, the map π^* makes $H_T^*(M)$ into a $\mathbb{C}[u_1, \dots, u_{\ell}]$ -module. Define $H_T^*(M)_{\text{loc}} = H_T^*(X) \otimes_{\mathbb{C}[u_1, \dots, u_{\ell}]} \mathbb{C}(u_1, \dots, u_{\ell})$. A fundamental result of the subject is the localization theorem:

Theorem 1. *Let $\{P\}$ denote the set of T -fixed components of M . Then $H_T^*(M)_{\text{loc}} \cong \bigoplus_P H^*(P) \otimes \mathbb{C}(u_1, \dots, u_{\ell})$.*

If P is a fixed component of M , the normal bundle to P splits as a sum over the characters of the T -action on the fibers: $N_P = \bigoplus_{\lambda} V_{\lambda}$. Let n_{λ}^i denote the formal Chern roots of V_{λ} . If we identify the equivariant parameters $u_1, \dots, u_{\ell}$ with linear forms on the Lie algebra of T , then the equivariant Euler class $e(P)$ of N_P is equal to $\prod_{\lambda} \prod_i (n_{\lambda}^i + u_{\lambda})$. Since none of the characters λ are equal to zero, we see that $e(P)$ is always invertible. In light of this fact, we can describe the above isomorphism more explicitly. The map $H_T^*(X)_{\text{loc}} \rightarrow \bigoplus_P H^*(P) \otimes \mathbb{C}(u_1, \dots, u_{\ell})$ is given by $\omega \mapsto \bigoplus_P \frac{i_P^* \omega}{e(P)}$, where $i_P : P \hookrightarrow X$ is the inclusion map.

If $f : M \rightarrow N$ is a proper map of T -spaces, we have the equivariant analogue of the cohomological push-forward $f_* : H_T^*(M) \rightarrow H_T^*(N)$. As in the non-equivariant setting, f_* satisfies the projection formula $f_*(f^*(\omega) \wedge \eta) = \omega \wedge f_* \eta$. The new feature in equivariant cohomology is that we have

an explicit expression for the restriction of $f_*\omega$ to a fixed component in N . This is given by the functorial localization formula [12] [13]:

Theorem 2. *Let $f : M \rightarrow N$ be a proper map of T -spaces. Let P be a fixed component of N and let $\{F\}$ be the collection of fixed components in M which f maps into P . Let $\omega \in H_T^*(M)$. Then:*

$$\sum_F f_* \frac{i_F^* \omega}{e(F)} = \frac{i_P^* f_* \omega}{e(P)}.$$

2.2. Power Series in Equivariant Cohomology. For simplicity, we assume that M has a $T = S^1$ action. Let X be the vectorfield on M induced by the action of T . Let $C(M)$ denote the ring of formal power series in u with coefficients in $\Omega^*(M)^T \otimes \mathbb{C}$. Then $d_X = d - ui_X : C(M) \rightarrow C(M)$. We define $H^*(C(M)) = \ker d_X / \text{im } d_X$.

Multiplication by u gives $H^*(C(M))$ the structure of a $\mathbb{C}[u]$ -module. Given any $\mathbb{C}[u]$ -module A , we define $A_{loc} = A \otimes \mathbb{C}(u)$. Here, we interpret rational polynomials of the form $\frac{1}{1+uf(u)}$ as convergent power series by expanding around $u = 0$.

Let g be a T -invariant metric on M , and define $\theta = g(X, \cdot)$. Let $F \subset M$ be the T -fixed locus, and let ρ be a bump function identically equal to one outside a tubular neighborhood of F , and equal to zero inside a smaller tubular neighborhood. (Note that we allow F to have different components of varying dimensions). Then $\rho(d_X \theta)^{-1}$ is a well-defined element of $C(M)_{loc}$. Furthermore, it is easy to see that for any closed form $\omega \in C(M)_{loc}$, $\omega - d_X(\omega \cdot \theta \cdot \rho(d_X \theta)^{-1})$ has compact support in a tubular neighborhood of F . It follows that every closed form in $C(M)_{loc}$ may be represented by a form with compact support in a tubular neighborhood of F . Let $C(\nu_F)_c$ denote the ring of formal power series in u whose coefficients are T -invariant forms with compact support in the normal bundle ν_F .

Let $\pi_* : C(\nu_F)_{c,loc} \rightarrow C(F)_{loc}$ be the map $\sum \omega_n u^n \mapsto \sum \pi_* \omega_n u^n$, where $\pi_* \omega_n$ denotes the integral of ω_n over the fiber of ν_F . We first remark that $\pi_* i_X \omega = 0$ for any differential form ω with compact support in ν_F . To see this, note that we can always express ω locally in the form $\omega = f(x, t) \pi^* \phi dt_1 \dots dt_k$. Here x represents the coordinates along F , t represents the coordinates along the fiber, and $\pi^* \phi$ is the pullback of a form on F via the projection $\pi : \nu_F \rightarrow F$. Thus $i_X \omega = f(x, t) \pi^* \phi \cdot i_X dt_1 \dots dt_k$ which clearly integrates to zero along the fiber, since the degree of the form along the fiber is necessarily smaller than the fiber dimension.

It follows that $\pi_* d_X = d\pi_*$, and therefore it induces a map in cohomology $\pi_* : H^*(C(\nu_F)_c)_{loc} \rightarrow H^*(C(F))_{loc}$.

Theorem 3. π_* is injective.

Proof. Let $\omega = \sum_n \omega_n u^n \in C(\nu_F)_{c,loc}$ be closed. Suppose $\pi_* \omega = d\eta$. By subtracting off $\pi^* d\eta \cdot \Phi$, where Φ is the equivariant Thom class of ν_F , we may reduce to the case in which $\pi_* \omega = 0$. For any differential form α , denote by $\alpha[i]$ the degree i part. Then for some $0 \leq k \leq \dim M$, $\omega = \omega[k] + \omega[k-1] + \dots + \omega[0]$, where $\omega[i] = \sum \omega_n[i] u^n$. Since ω is d_X -closed, $d\omega_n[k] = 0$. Since $\pi_* \omega[k] = \sum \pi_* \omega_n[k] u^n = 0$, by the Thom isomorphism theorem, we can find compactly supported forms y_n such that $dy_n = \omega_n[k]$. By averaging over T , we may assume that these forms are T -invariant. Then $\omega - d_X \sum y_n u^{n-1}$ has top degree $< k$ and is annihilated by π_* . The proof then follows by induction. $\square$

Since $[\omega] = 0$ if and only if $\pi_*[\omega] = 0$, we see that $[\omega] = [\pi^* \pi_* \omega \cdot \Phi]$. Hence $[\omega]|_F = [\pi_* \omega \cdot e(\nu_F)]$, where $e(\nu_F)$ is the equivariant Euler class. Since the Euler class is invertible, the restriction map $[\omega] \mapsto [\omega]|_F$ must be invertible. This proves the localization theorem for $H^*(C(M))_{loc}$:

Theorem 4. *Let $F \subset M$ denote the fixed locus of T . Then the restriction map $H^*(C(M))_{loc} \rightarrow H^*(C(F))_{loc}$ is an isomorphism.*

We now introduce the ring of analytic forms: Let $C^{an}(M) \subset C(M)$ denote the ring of forms $\sum \omega_n u^n$ with the property that the partial sums $\sum_{n=0}^N \omega_n s^n$ converge in the C^∞ sense to a form $\omega \in \Omega^*(M)^T$ for $\|s\|$ sufficiently small. Let $B^{an}(M) = d_X(C(M)) \cap C^{an}(M)$. We define

$$H^*(C^{an}(M)) = \frac{\ker d_X : C^{an}(M) \rightarrow C^{an}(M)}{B^{an}(M)}.$$

If F denotes the T -fixed locus of M , the above proof of the localization theorem extends word for word to the case of analytic forms.

For $s \in \mathbb{C}^*$, following Witten [18], let $d_s = d - si_X : \Omega^*(M)^T \rightarrow \Omega^*(M)^T$. Define $H_s^*(M) = \frac{\ker d_s}{\text{im } d_s}$. Again, the above proof of the localization theorem adapts easily to the case of $H_s^*(M)$. Define an equivalence relation $\sim$ on $\prod_{s \in \mathbb{C}^*} H_s^*(M)$ as follows: For $\omega, \omega' \in \prod_{s \in \mathbb{C}^*} H_s^*(M)$, we say that $\omega \sim \omega'$ if $\omega_s = \omega'_s$ for $\|s\|$ sufficiently small. By ω_s , of course, we mean the s -component of ω . We denote the group $\prod_{s \in \mathbb{C}^*} H_s^*(M) / \sim$ by W .

Proposition 1. *There is a natural evaluation map $ev : H^*(C^{an}(M)) \rightarrow W$ given by $[\sum \omega_n u^n] \mapsto [\sum \omega_n s^n]$.*

Proof. Let $\omega = \sum \omega_n u^n \in C^{an}(M)$ be a closed form, and let $d_X \sum \eta_n u^n \in C^{an}(M)$. We wish to show that for $\|s\|$ sufficiently small, $[\sum \omega_n s^n] = [\sum \omega_n s^n + d_s \sum \eta_n s^n]$ in $H_s^*(M)$. First, for $\|s\|$ sufficiently small, both sums $\sum \omega_n s^n$ and $\sum d_s \eta_n s^n$ converge. Therefore, it suffices to prove that $\sum d_s \eta_n s^n$ is exact for these values of s . Let $F \subset M$ be the T -fixed locus. Then $\sum d_s \eta_n|_F s^n = \sum d\eta_n|_F s^n$. By Hodge theory, $\sum d\eta_n|_F s^n$ converges to

an exact form. Hence $[\sum d_s \eta_n s^n]|_F = 0$, and therefore $[\sum d_s \eta_n s^n] = 0$ by localization. $\square$

Let $F(x_1, \dots, x_r)$ be function which is holomorphic in a neighborhood of $(0, \dots, 0)$. Let $F = \sum_{i_1, \dots, i_r} a_{i_1, \dots, i_r} x_1^{i_1} \cdots x_r^{i_r}$ be the corresponding power series expansion. Let $[\omega_1], \dots, [\omega_r]$ be equivariant forms in $H_T^*(M)$ such that $u | \deg_0 \omega_i$. Note that this property is independent of the choice of representatives for $[\omega_i]$. Then $F([\omega_1], \dots, [\omega_r])$ is a well-defined element of $H^*(C^{an}(M))$. We can see this as follows:

Fix representatives $\omega_1, \dots, \omega_r \in C^{an}(M)$. Write $\omega_i = \tilde{\omega}_i + f_i u$, where $f_i \in C^\infty(M) \otimes \mathbb{C}[u]$ and $\deg \tilde{\omega}_i > 0$. Then

$$\begin{aligned} F(\omega_1, \dots, \omega_r) &= \sum_{i_1, \dots, i_r} a_{i_1, \dots, i_r} (\tilde{\omega}_1 + f_1 u)^{i_1} \cdots (\tilde{\omega}_r + f_r u)^{i_r} \\ &= \sum_{j_1, \dots, j_r} \tilde{\omega}_1^{j_1} \cdots \tilde{\omega}_r^{j_r} \sum_{i_1, \dots, i_r} a_{i_1, \dots, i_r} \binom{i_1}{j_1} \cdots \binom{i_r}{j_r} (f_1 u)^{i_1 - j_1} \cdots (f_r u)^{i_r - j_r} \\ &= \sum_{j_1, \dots, j_r} \partial^{(j_1)} \cdots \partial^{(j_r)} F(f_1 u, \dots, f_r u) \tilde{\omega}_1^{j_1} \cdots \tilde{\omega}_r^{j_r}. \end{aligned}$$

Here $\partial^{(k)} = \frac{\partial^k}{k!}$. The sum over $j_1, \dots, j_r \geq 0$ is finite because the forms $\tilde{\omega}_i$ are nilpotent. It is therefore clear that the above expression is a well-defined d_X -closed form in $C^{an}(M)$. We next show that the corresponding class in $H^*(C^{an}(M))$ is independent of the choice of generators for $[\omega_i]$. Let $H(s_1, t_1, \dots, s_r, t_r) = F(s_1 + t_1, \dots, s_r + t_r)$. Then $H = F(s_1, \dots, s_r) + t_1 \tilde{H}_1 + \dots + t_r \tilde{H}_r$. Thus, $F(\omega_1 + d_X \eta_1, \dots, \omega_r + d_X \eta_r) = H(\omega_1, d_X \eta_1, \dots, \omega_r, d_X \eta_r) = F(\omega_1, \dots, \omega_r) + d_X(\eta_1 \tilde{H}_1 + \dots + \eta_r \tilde{H}_r)$, so $F([\omega_1], \dots, [\omega_r])$ is well-defined independent of our choice of generators. Before proceeding further, we point out an important property possessed by the forms of this type.

Let $\hat{C}(M) \subset C^{an}(M)$ denote the subring $\{\text{closed invariant forms} \otimes \mathbb{C}\{u\}\}$ where $\mathbb{C}\{u\}$ denotes the ring of power series in u which converge in sufficiently small neighborhoods of the origin. Let $\hat{H}(M) \subset H^*(C^{an}(M))_{loc}$ denote the subspace of forms $[\omega]$ such that $\omega|_F \in \hat{C}(F)_{loc}$ for some representative $\omega \in [\omega]$. Then if $[\omega_1], \dots, [\omega_r] \in H_T^*(M)$, and $u | \deg_0 \omega_i$, then $F([\omega_1], \dots, [\omega_r]) \in \hat{H}(M)$. This is because $\deg_0 \omega_i|_F \in \mathbb{C}[u]$ and $d\tilde{\omega}_i|_F = 0$. Note also that functorial localization holds in this situation, and implies that, for $f : M \rightarrow N$, $f_* \hat{H}(M) \subset \hat{H}(N)$.

Lemma 1. *ev : $\hat{H}(M) \rightarrow W(M)$ is injective.*

Proof. Let $[\omega] \in \hat{H}(M)$, so that $\omega|_F = \omega_1 f_1 + \dots + \omega_k f_k$, where ω_i are closed forms on F and f_i are convergent Laurent series in u in some small disk

about the origin. If $[\omega] \neq 0$, then by localization, $[\omega]|_F \neq 0$. Without loss of generality, we may assume that $\omega_1, \dots, \omega_k$ are linearly independent as cohomology classes. We can always choose an arbitrarily small s so that $f_1(s), \dots, f_k(s)$ are not all zero. We then have that $ev[\omega]_s|_F = f_1(s)[\omega_1] + \dots + f_k(s)[\omega_k] \neq 0$. Hence $ev[\omega] \neq 0$ in $W(M)$. $\square$

All the above machinery was put together to make the following argument: Let $f : M \rightarrow N$, and let $F = F([\omega_1], \dots, [\omega_r])$ for $[\omega_i] \in H_T^*(M)$, $u | \deg_0 \omega_i$. Suppose $\theta_n(s)$ is a sequence of d_s -closed forms which converge in the C^∞ sense to a representative of $(evF)_s$ for $\|s\|$ sufficiently small. If we factor f as an inclusion followed by a projection, then f_* makes sense on the form level and $f_*\theta_n(s)$ converge to representatives of $(f_*evF)_s$. Now suppose that $[f_*\theta_n(s)] = [b_n(s)]$ and that $b_n(s) \rightarrow b(s)$. Then $f_*\theta_n(s) = b_n(s) + d_s\eta_n(s)$, and since $f_*\theta_n(s)$ and $b_n(s)$ converge, we have that $[b(s)] = [f_*evF]_s$. Now if $s \mapsto [b(s)]$ corresponds to a form evG for some $G([\alpha_1], \dots, [\alpha_k]), [\alpha_i] \in H_T^*(N)$, then $f_*evF = evf_*F = evG$, and therefore $f_*F = G$. Thus, we can compute push-forwards of summations by applying the push-forward term-by-term, so long as the corresponding summation converges in the sense discussed in this section. This observation will be used implicitly throughout this paper; for example, it will allow us in sections 5 and 9 to reduce computations involving convergent power series to computations involving polynomials.

Remark 1. In what follows we will be interested in the case where T is a compact torus of arbitrary dimension. It is easy to see how to generalize the above machinery to this situation. The difficulty lies more in the notation than in any other aspect.

3. THE ORBIFOLD ELLIPTIC CLASS

Let X be a smooth projective variety with a holomorphic $T \times G$ action. Let $D = \sum_I \alpha_i D_i$ be a smooth G -normal crossing divisor with $\alpha_i < 1$. The G -normal condition means that if $g \in \text{stab}_G(x)$ and $x \in D_i$, then $g \cdot D_i = D_i$. Let $X_\gamma^{g,h}$ be a connected component of the fixed locus $X^{g,h}$ for some commuting pair $g, h \in G$. Then $N_{X_\gamma^{g,h}/X}$ splits into character subbundles $\bigoplus_\lambda N_\lambda$, where g (resp. h) acts on N_λ as multiplication by $e^{2\pi i \lambda(g)}$ (resp. $e^{2\pi i \lambda(h)}$). Let $I_\gamma^{g,h} \subset I$ index the set of divisors D_i^X which contain $X_\gamma^{g,h}$. Since D is G -normal, g (resp. h) acts on $\mathcal{O}(D_i)|_{X_\gamma^{g,h}}$ as multiplication by $e^{2\pi i \lambda_i(g)}$ (resp. $e^{2\pi i \lambda_i(h)}$) for every $i \in I_\gamma^{g,h}$. For $i \notin I_\gamma^{g,h}$, we define $\lambda_i(g) = \lambda_i(h) = 0$.

Following [7], we define the orbifold elliptic class associated to the pair $(X_\gamma^{g,h}, D)$ as follows: $\mathcal{E}ll_{orb}(X_\gamma^{g,h}, D) =$

$$(i_{X_\gamma^{g,h}})_* \left\{ \prod_{TX_\gamma^{g,h}} \frac{\frac{x_i}{2\pi i} \theta(\frac{x_i}{2\pi i} - z) \theta'(0)}{\theta(\frac{x_i}{2\pi i}) \theta(-z)} \prod_{\lambda, N_\lambda} \frac{\theta(\frac{x_{\lambda,c}}{2\pi i} + \lambda(g) - \lambda(h)\tau - z) \frac{\theta'(0)}{2\pi i}}{\theta(\frac{x_{\lambda,c}}{2\pi i} + \lambda(g) - \lambda(h)\tau) \theta(-z)} \right\} \\ \prod_I \frac{\theta(\frac{D_i}{2\pi i} + \lambda_i(g) - \lambda_i(h)\tau - (-\alpha_i + 1)z) \theta(-z)}{\theta(\frac{D_i}{2\pi i} + \lambda_i(g) - \lambda_i(h)\tau - z) \theta(-(-\alpha_i + 1)z)}$$

Here x_i denote the equivariant Chern roots of $TX_\gamma^{g,h}$, $x_{\lambda,c}$ denote the equivariant Chern roots of N_λ , and (abusing notation) D_i denote the equivariant first Chern classes of the corresponding divisors.

Finally we define the orbifold elliptic class of (X, D, G) as follows:

$$\mathcal{E}ll_{orb}(X, D, G) = \frac{1}{|G|} \sum_{gh=hg, \gamma} \mathcal{E}ll_{orb}(X_\gamma^{g,h}, D).$$

When $G = 1$, we will write $\mathcal{E}ll(X, D)$ instead of $\mathcal{E}ll_{orb}(X, D, G)$ and refer to this object as the equivariant elliptic class of the pair (X, D) . We view all such classes as an elements inside the ring $\widehat{H}(X)$.

Let $\{F\}$ denote the collection of fixed components of X . For each F let $e(F)$ denote the equivariant Euler class of the normal bundle to F . We define the equivariant orbifold elliptic index

$$Ell_{orb}(X, D, G) \equiv \left(\frac{2\pi i \theta(-z)}{\theta'(0)} \right)^{\dim X} \sum_F \int_F \frac{\mathcal{E}ll_{orb}(X, D, G)}{e(F)}.$$

It is a convergent power series in the equivariant parameters which depends implicitly upon the value of the complex parameter z and on the lattice parameter τ used in the definition of the Jacobi theta function. We define the equivariant elliptic index $Ell(X, D)$ similarly.

4. TORIC VARIETIES AND EQUIVARIANT COHOMOLOGY

For a good reference on toric varieties, see [10]. Let X be a smooth complete toric variety of dimension n . We denote the fan of X by Σ_X , the lattice of X by N_X , and the big torus by T_X . Let Y be a smooth complete toric variety which satisfies the following properties:

- (1): $N_X \subset N_Y$ is a finite index sublattice.
- (2): Σ_X is a refinement of Σ_Y obtained by adding finitely-many one dimensional rays.

There is an obvious map of fans $\nu : \Sigma_X \rightarrow \Sigma_Y$ which induces a smooth map $\mu : X \rightarrow Y$. We call a map induced by such a morphism of fans a *toric morphism*. It is easy to verify that $\mu : T_X \rightarrow T_Y$ is a covering map with covering group N_Y/N_X . Thus, we may regard Y as a T_X -space. Our

goal in this section is to obtain a convenient description of the equivariant pushforward $\mu_* : H_T^*(X) \rightarrow H_T^*(Y)$ in terms of the combinatorics of Σ_X and Σ_Y . Here $T = T_X$.

We first note that fixed points F of X are in 1 – 1 correspondence with n -dimensional cones $C_F \subset \Sigma_X$. Furthermore, the infinitesimal weights of the T -action on N_F correspond to linear forms in $\text{Hom}(N_X, \mathbb{Z})$ which are dual to the generators of C_F in N_X . With this identification in mind, for any $\omega \in H_T^*(X)$, the collection of polynomials $\{\omega|_F\}_{F \in \text{Fix}(X)}$ defines a piecewise polynomial function on the fan Σ_X . Define $\mathbb{C}[\Sigma_X]$ to be the ring of all piecewise polynomial functions on the fan of X . It is well-known that the map $H_T^*(X) \rightarrow \mathbb{C}[\Sigma_X]$ described here is an isomorphism:

Theorem 5. $H_T^*(X) \cong \mathbb{C}[\Sigma_X]$.

Via this identification, we define $\nu_* : \mathbb{C}[\Sigma_X] \rightarrow \mathbb{C}[\Sigma_Y]$ to be the map which makes the following diagram commute:

$$\begin{array}{ccc} \mathbb{C}[\Sigma_X] & \xrightarrow{\nu_*} & \mathbb{C}[\Sigma_Y] \\ \parallel & & \parallel \\ H_T^*(X) & \xrightarrow{\mu_*} & H_T^*(Y) \end{array}$$

Here we understand $\mathbb{C}[\Sigma_Y]$ to be the ring of piecewise polynomial functions on Σ_Y with respect to the lattice N_X .

We now describe ν_* more explicitly. First notice that for $f \in \mathbb{C}[\Sigma_X]$, $\nu_* f$ is given by viewing $f|_F$ as the zero degree part of an equivariant cohomology class $\omega \in H_T^*(X)$, pushing ω forward by μ_* , and then forming the piecewise polynomial function defined by the zero degree part of $\mu_* \omega$. Thus, let $C \subset \Sigma_Y$ be an n -dimensional cone. Let $\nu^{-1}C$ be the fan $\Sigma_C \subset \Sigma_X$ which is the union of n -dimensional cones C_i . Let $x_1^{C_i}, \dots, x_n^{C_i}$ be the linear forms dual to C_i and $x_1^C, \dots, x_n^C$ the linear forms in $\text{Hom}(N_Y, \mathbb{Z}) \subset \text{Hom}(N_X, \mathbb{Z})$ dual to C . By functorial localization:

$$(\nu_* f)_C = \sum_{C_i \subset \Sigma_X} f_{C_i} \frac{\prod_{j=1}^n x_j^C}{\prod_{j=1}^n x_j^{C_i}}.$$

Similarly, we define $\nu^* : \mathbb{C}[\Sigma_Y] \rightarrow \mathbb{C}[\Sigma_X]$ to be the map which makes the following diagram commute:

$$\begin{array}{ccc} \mathbb{C}[\Sigma_Y] & \xrightarrow{\nu^*} & \mathbb{C}[\Sigma_X] \\ \parallel & & \parallel \\ H_T^*(Y) & \xrightarrow{\mu^*} & H_T^*(X) \end{array}$$

Proposition 2. $\nu^*(f) = f \circ \nu$

Proof. Let $\omega \in H_T^*(Y)$ be the form such that $\omega|_P = f|_P$ for every fixed point P . Let $F \in \mu^{-1}(P)$. Then

$$\begin{array}{ccc} H_T^*(Y) & \xrightarrow{\mu^*} & H_T^*(X) \\ \downarrow & & \downarrow \\ H_T^*(P) & \xrightarrow{\mu_F^*} & H_T^*(F) \end{array}$$

commutes. Hence $(\mu^*\omega)|_F = \mu_F^*(\omega|_P) = \mu_F^*(f_P) = f_P$. Thus $\nu^*(f)$ is the piecewise polynomial function which is equal to f_{C_P} on every cone $C_F \in \nu^{-1}C_P$. This is precisely the piecewise polynomial $f \circ \nu$. $\square$

The map $\nu^* : \mathbb{C}[\Sigma_Y] \rightarrow \mathbb{C}[\Sigma_X]$ makes $\mathbb{C}[\Sigma_X]$ into a $\mathbb{C}[\Sigma_Y]$ -module. As such, we observe:

Proposition 3. ν_* is a $\mathbb{C}[\Sigma_Y]$ -module homomorphism.

Proof. In other words, we wish to prove the projection formula $\nu_*(f\nu^*g) = \nu_*(f) \cdot g$. This follows from identifying ν_* with μ_* , ν^* with μ^* and invoking the projection formula from equivariant cohomology. $\square$

5. TOROIDAL EMBEDDINGS AND TOROIDAL MORPHISMS

5.1. Definitions. Let X be a compact variety and $D_X = \sum_{I \in I_X} D_i^X$ a divisor on X whose irreducible components are smooth normal crossing divisors. For $I \subset I_X$, let $X_{I,j}$ denote the j th connected component of $\cap_I D_i^X$. Let $X_{I,j}^o = X_{I,j} - \cup_{I' \subsetneq I} X_{I',j}^o$. The collection of subvarieties $X_{I,j}^o$ form a stratification of X . Associated to these data is a polyhedral complex with integral structure defined as follows:

Corresponding to $X_{I,j}$, define $N_{I,j} = \mathbb{Z}e_{i_1,j} + \dots + \mathbb{Z}e_{i_k,j}$ to be the free group on the elements $e_{i_1,j}, \dots, e_{i_k,j}$. Here $i_1, \dots, i_k$ are the elements of I . Define $C_{I,j}$ to be the cone in the first orthant of this lattice. Whenever $I' \subset I$ and $X_{I,j} \subset X_{I',j'}$ we have natural inclusion maps $N_{I',j'} \hookrightarrow N_{I,j}$ and $C_{I',j'} \hookrightarrow C_{I,j}$. Define Σ_X to be the polyhedral complex with integral structure obtained by gluing the cones $C_{I,j}$ together according to these inclusion maps.

Let $\mathbb{C}[\Sigma_X]$ denote the ring of piecewise polynomial functions on Σ_X . Fix $C \subset \Sigma_X$. Define f^C to be the piecewise polynomial function which is equal to $\prod_{j=1}^{\dim C} x_j^C$ on every cone containing C , and equal to zero everywhere else. As in the toric geometry case, there is a natural correspondence between

piecewise linear functions on Σ_X and Cartier divisors whose irreducible components are components of D_X . We denote the piecewise linear function corresponding to D by f^D .

5.2. Toroidal Morphisms. Our primary interest in this section is the study of toroidal morphisms. This is a map $\mu : (X, D_X, \Sigma_X) \rightarrow (Y, D_Y, \Sigma_Y)$ which satisfies the following:

- (1): $\mu : X - D_X \rightarrow Y - D_Y$ is an unramified cover.
- (2): μ maps the closure of a stratum in X to the closure of a stratum in Y .
- (3): Let U_y be an analytic neighborhood of $y \in Y$ such that the components of D_Y passing through y correspond to coordinate hyperplanes. Then for $x \in \mu^{-1}(y)$, there exists an analytic neighborhood U_x of x such that the components of D_X passing through x correspond to coordinate hyperplanes of U_x . Moreover, the map $U_x \rightarrow U_y$ is given by monomial functions in the coordinates.

Corresponding to μ , we can define a map $\nu : \Sigma_X \rightarrow \Sigma_Y$ as follows: Let $C_{I,i} \subset \Sigma_X$ and let $e_1, \dots, e_k \in N_{I,i}$ be the generators of $C_{I,i}$ which correspond to the divisors $D_1^X, \dots, D_k^X$. We have that $\mu(X_{I,i}) = Y_{J,j}$. Let $v_1, \dots, v_\ell \in N_{J,j}$ be the generators of $C_{J,j}$ which correspond to the divisors $D_1^Y, \dots, D_\ell^Y$. For $1 \leq s \leq k$, $1 \leq t \leq \ell$, define a_{st} to be the coefficient of D_s^X of the divisor $\mu^*(D_t^Y)$. Then we define $\nu(e_s) = \sum a_{st} v_t$. Note that if $(X, \Sigma_X) \rightarrow (Y, \Sigma_Y)$ is a smooth toric morphism of toric varieties, then $\nu : \Sigma_X \rightarrow \Sigma_Y$ is the natural morphism of polyhedral complexes.

We have the following proposition relating ν to μ :

Proposition 4. *If $C = C_{J,j} \subset \Sigma_Y$, then $\nu^{-1}C$ is the union of fans $\Sigma_\alpha \subset \Sigma_X$ with the following properties:*

- (1): Σ_α is a refinement of C obtained by adding finitely-many 1-dim rays.
- (2): The lattice N_α of Σ_α is a finite index sub-lattice of N_C .
- (3): The fans Σ_α are in 1-1 correspondence with connected components U_α of $\mu^{-1}(N_{Y_{J,j}}^\circ)$. The map $U_\alpha \rightarrow N_{Y_{J,j}}^\circ$ is a fibration given by the smooth toric morphism $\mathbf{P}_{\Sigma_\alpha, N_\alpha} \rightarrow \mathbf{P}_{C, N_C}$ along the fiber, and a $d_\alpha = d(\Sigma_\alpha)$ -cover of $Y_{J,j}^\circ$ along the base.

For a proof, see [7].

5.3. Pushforward formula for Polyhedral Complexes. Motivated by the description of the push-forward ν_* for toric morphisms, define $\nu_* : \mathbb{C}[\Sigma_X] \rightarrow \mathbb{C}[\Sigma_Y]$ as follows. Let $C \subset \Sigma_Y$ be an n -dimensional cone with dual linear forms $x_1^C, \dots, x_n^C$. Then for $f \in \mathbb{C}[\Sigma_X]$, we define:

$$(\nu_* f)_C = \sum_{\alpha} d_{\alpha} \sum_{C_i \in \Sigma_{\alpha}} f_{C_i} \cdot \frac{\prod_{j=1}^n x_j^C}{\prod_{j=1}^n x_j^{C_i}}$$

The second sum is taken over the cones $C_i \subset \Sigma_{\alpha}$ with the same dimension as C .

Let V be the toric variety $\coprod_{\alpha} d_{\alpha} \cdot \mathbf{P}_{\Sigma_{\alpha}, N_{\alpha}}$ with polyhedral fan Σ_V . We have a natural toric morphism $V \rightarrow \mathbb{C}^n$. We can compactify V and $\mathbb{C}^n$ to obtain a smooth toric morphism $\overline{V} \rightarrow \mathbb{P}^n$. If we view f as a piece-wise polynomial function on the fan of $\overline{V}$, then the above formula simply corresponds to $(\nu_* f)_C$ where $\nu : \Sigma_{\overline{V}} \rightarrow \Sigma_{\mathbb{P}^n}$. This identification allows us to apply the tools of the previous section toward the study of ν_* .

We first observe that $(\nu_* f)_C$ is indeed a polynomial function. This follows from the above identification of ν_* with the equivariant pushforward of a toric morphism. Furthermore, if we define $\nu^* : \mathbb{C}[\Sigma_Y] \rightarrow \mathbb{C}[\Sigma_X]$ by the formula $\nu^*(f) = f \circ \nu$ then the projection formula:

$$\nu_*(f \nu^* g)_C = \nu_*(f)_C \cdot g_C$$

follows from the projection formula in equivariant cohomology.

Proposition 5. $\nu_*(f)$ is a piece-wise polynomial function.

Proof. We first show that $\nu_*(f^C)$ is piece-wise polynomial.

Fix $f = f^C$. Suppose $\nu(C) \subset C_0$ for some $C_0 \subset \Sigma_Y$ of dimension $k = \dim C$. Then $\nu_*(f)_{C_0} = d(\Sigma_{C_0}) \prod_{j=1}^k x_j^{C_0}$. Suppose C_1 is a cone containing C_0 . We wish to show $(\nu_* f)_{C_1}$ is an extension of $(\nu_* f)_{C_0}$.

Consider the toric morphism $\sigma : \mathbf{P}_{\Sigma_{C_1}, N(\Sigma_{C_1})} \rightarrow \mathbb{C}^{\dim C_1}$ induced by the map $\nu : \Sigma_{C_1} \rightarrow C_1$. Let $D_1, \dots, D_k$ be the divisors in $\mathbf{P}_{\Sigma_{C_1}, N(\Sigma_{C_1})}$ which correspond to the generators of C . Then the piece-wise polynomial function $f \in \mathbb{C}[\Sigma_{C_1}]$ represents the equivariant Thom class of $D_1 \cap \dots \cap D_k$. Since $\sigma(D_1 \cap \dots \cap D_k)$ is the affine subspace of $\mathbb{C}^{\dim C_1}$ corresponding to C_0 , we have that $\sigma_*(f)$ is the degree of σ along $D_1 \cap \dots \cap D_k$ times the polynomial function which represents the equivariant Thom class of this subspace. But this implies that:

$$\nu_*(f)_{C_1} = d(\Sigma_{C_1}) \frac{[N(\Sigma_{C_1}) : N(C_1)]}{[N(\Sigma_{C_0}) : N(C_0)]} \prod_{j=1}^k x_j^{C_0} = d(\Sigma_{C_0}) \prod_{j=1}^k x_j^{C_0}.$$

We need to explain the last equality. If C_0 corresponds to the strata $Y_{I,j}^o$ and $U \rightarrow N_{Y_{I,j}^o}$ is the fibration in Proposition 4 corresponding to the subdivision Σ_{C_0} , then $d(\Sigma_{C_0})[N(\Sigma_{C_0}) : N(C_0)]$ and $d(\Sigma_{C_1})[N(\Sigma_{C_1}) : N(C_1)]$ both give the number of points in the pre-image of a generic point in $N_{Y_{I,j}^o}$.

Next suppose that C is mapped to a cone C_0 of strictly larger dimension. Consider the toric morphism $\mathbf{P}_{\Sigma_{C_0}, N(\Sigma_{C_0})} \rightarrow \mathbf{P}_{C_0, N(C_0)}$ induced by the map $\nu : \Sigma_{C_0} \rightarrow C_0$. The polynomial function $f \in \mathbb{C}[\Sigma_{C_0}]$ represents the Thom class of an exceptional toric subvariety. Thus $\nu_*(f) = 0$, and it is easy to verify that $\nu_*(f) = 0$ on every cone containing C_0 . Thus, ν_* maps the elements f^C to piecewise polynomial functions. Since these functions generate $\mathbb{C}[\Sigma_X]$ as a $\mathbb{C}[\Sigma_Y]$ -module, the proposition follows from the projection formula. $\square$

In what follows we assume that $\mu : X \rightarrow Y$ is an equivariant map of projective T -spaces. Furthermore, we assume that the irreducible components of D_X and D_Y are invariant under the T -action. Define a map $\rho_X : \mathbb{C}[\Sigma_X] \rightarrow H_T^*(X)$ as follows: Fix a cone $C = C_{I,i}$ which corresponds to a connected component of the intersection locus of the divisors $D_1, \dots, D_k$. Define $\rho_X[f^C \cdot (f^{D_1})^{a_1} \dots (f^{D_k})^{a_k}] = \Phi_{X_{I,i}} \wedge D_1^{a_1} \wedge \dots \wedge D_k^{a_k}$. Here $\Phi_{X_{I,i}}$ denotes the equivariant Thom class of $X_{I,i} \subset X$ and, by abuse of notation, D_j denote the equivariant Thom classes of the divisors D_j .

Lemma 2. ρ_X is a ring homomorphism.

Proof. Fix cones $C_1 = C_{I_1, i_1}$ and $C_2 = C_{I_2, i_2}$. It suffices to prove the theorem for the polynomials f^{C_1} and f^{C_2} . Let $I = I_1 \cup I_2$. Let $C_{I,i}$ denote the cones which correspond to components of the intersection $X_{I_1, i_1} \cap X_{I_2, i_2}$. Clearly

$$f^{C_1} f^{C_2} = \sum_{I,i} f^{C_{I,i}} \prod_{I_1 \cap I_2} f^{D_j}.$$

Thus $\rho_X(f^{C_1} f^{C_2}) = \sum_{I,i} \Phi_{X_{I,i}} \prod_{I_1 \cap I_2} D_j$. However, by the equivariant version of the excess intersection formula, this is precisely the formula for $\rho_X(f^{C_1}) \rho_X(f^{C_2})$. $\square$

Lemma 3. $\rho_X \nu^* = \mu^* \rho_Y$.

Proof. It suffices to check this for polynomials $f^{C_{I,k}}$. If D is a divisor on Y whose irreducible components are components of D_Y , then $\nu^* f^D$ is the piecewise linear function corresponding to $\mu^* D$. It follows that $\rho_X \nu^* f^D = \mu^* \rho_Y f^D$. Since all the maps are ring homomorphisms, this implies that $\rho_X \nu^* \prod_{j \in I} f^{D_j} = \mu^* \rho_Y \prod_{j \in I} f^{D_j}$. Let $\mu^* D_i = \sum_j a_{ij} E_j$ as Cartier divisors. As in the lemma in the Appendix, choose equivariant Thom forms Φ_{E_j} and Φ_{D_i} with support in small tubular neighborhoods of their respective divisors so that:

$$\mu^* \Phi_{D_i} = \sum_j a_{ij} \Phi_{E_j} + d\psi_i$$

as forms. Here ψ_i are equivariant forms with compact support in $\mu^{-1}N_{D_i}$. Let $\{I, k\}$ index the connected components of $\cap_I D_i$. If we choose N_{D_i} sufficiently small, then

$$\prod_I \Phi_{D_i} = \sum_{I,k} (\prod_I \Phi_{D_i})_{I,k}$$

where $(\prod_I \Phi_{D_i})_{I,k}$ is the extension by zero of the form $\prod_I \Phi_{D_i}|_{N_{I,k}}$.

Now $\prod_I f^{D_i} = \sum f^{C_{I,k}}$ and clearly $(\prod_I \Phi_{D_i})_{I,k}$ is a representative of $\rho_Y(f^{C_{I,k}})$. We have that

$$\mu^*(\prod_I \Phi_{D_i})_{I,k} = \{ \prod_I (\sum_j a_{ij} \Phi_{E_j} + d\psi_i) \}_{\mu^{-1}N_{I,k}}$$

where the subscript $\mu^{-1}N_{I,k}$ means the extension by zero of the form restricted to this open set. Since the ψ_i forms have compact support in $\mu^{-1}N_{D_i}$, this form is cohomologous to

$$\{ \prod_I \sum_j a_{ij} \Phi_{E_j} \}_{\mu^{-1}N_{I,k}}.$$

But this is in turn a representative of $\rho_X \nu^* f^{C_{I,k}}$. $\square$

Lemma 4. $\mu_* \rho_X = \rho_Y \nu_*$.

Proof. Since $\rho_X \nu^* = \mu^* \rho_Y$ and the polynomials f^C generate $\mathbb{C}[\Sigma_X]$ as a $\mathbb{C}[\Sigma_Y]$ -module, by the projection formula it suffices to check $\mu_* \rho_X f^C = \rho_Y \nu_* f^C$.

Case 1: $C_{I,i}$ is mapped by ν to a cone $C_{J,j}$ of the same dimension.

From the proof of Proposition 4, $\nu_* f^{C_{I,i}} = df^{C_{J,j}}$ where d is the degree of $\mu : X_{I,i} \rightarrow Y_{J,j}$. Thus, $\rho_Y \nu_* f^{C_{I,i}} = d\Phi_{Y_{J,j}} = \mu_* \nu_* f^{C_{I,i}}$.

Case 2: $C_{I,i}$ is mapped by ν into a cone of strictly larger dimension.

As shown in Proposition 4, $\nu_* f^{C_{I,i}} = 0$, so $\rho_Y \nu_* f^{C_{I,i}} = 0 = \mu_* \Phi_{X_{I,i}} = \mu_* \rho_X f^{C_{I,i}}$. $\square$

Remark 2. It is clear using the formalism of section 2 that the above lemmas relating μ to ν extend without difficulty to the ring $\mathbb{C}[[\Sigma_X]]$ of piecewise convergent power series. In this situation, ρ_X is a map from $\mathbb{C}[[\Sigma_X]]$ into $\hat{H}(X)$.

6. ELLIPTIC GENERA AND TOROIDAL MORPHISMS

Let $(\hat{X}, \sum_{I_{\hat{X}}} \hat{D}_j)$ and $(X, \sum_{I_X} D_i)$ be smooth projective toroidal embeddings with T -actions which leave $\hat{D}_j$ and D_i invariant. Let G be a finite

group which acts toroidally on $\hat{X}$ and commutes with the action of G . Suppose that $\mu : \hat{X} \rightarrow X$ is a T -equivariant toroidal morphism which is birational to a global quotient by G . If α_i are coefficients of the irreducible components D_i , define β_j so that $\mu^*(K_X + \sum \alpha_i D_i) = K_{\hat{X}} + \sum \beta_j \hat{D}_j$. Then:

Theorem 6.

$$\mu_* \mathcal{E}ll_{orb}(\hat{X}, \sum \beta_j \hat{D}_j, G) = \mathcal{E}ll(X, \sum \alpha_i D_i).$$

Proof. The proof of this theorem follows almost word for word the proof of Theorem 5.1 in [7]. The only difference is that here we examine the equivariant push-forward of equivariant cohomology classes, whereas [7] examine the push-forward of (non-equivariant) classes in the Chow ring. We reproduce the proof here for completeness.

We refer frequently to the notation in the previous section: For commuting elements $g, h \in G$, let $\hat{X}_{\gamma}^{g,h}$ denote the γ -th fixed component of (g, h) . Since the action of G on $\hat{X}$ is toroidal, $\hat{X}^{g,h}$ may be identified with $X_{I^{g,h},i}$ for some indexing set $I^{g,h} \subset I_{\hat{X}}$.

Consider the following class in $\hat{H}(\hat{X})$:

$$\begin{aligned} E = & \frac{1}{|G|} \sum_{gh=hg; \hat{X}_{\gamma}^{g,h}} \Phi_{\hat{X}_{\gamma}^{g,h}}^T \cdot \prod_{I_{\hat{X}} - I_{\gamma}^{g,h}} \frac{\frac{\hat{D}_j}{2\pi i} \theta(\frac{\hat{D}_j}{2\pi i} - (-\beta_j + 1)z) \theta'(0)}{\theta(\frac{\hat{D}_j}{2\pi i}) \theta(-(-\beta_j + 1)z)} \\ & \prod_{I_{\gamma}^{g,h}} \frac{\theta(\frac{\hat{D}_j}{2\pi i} + \epsilon_j(g) - \epsilon_j(h)\tau - (-\beta_j + 1)z) \frac{\theta'(0)}{2\pi i}}{\theta(\frac{\hat{D}_j}{2\pi i} + \epsilon_j(g) - \epsilon_j(h)\tau) \theta(-(-\beta_j + 1)z)} e^{2\pi i(-\beta_j + 1)\epsilon_j(h)z}. \end{aligned}$$

Here $\Phi_{\hat{X}_{\gamma}^{g,h}}^T$ is the equivariant Thom class of the subvariety $\hat{X}_{\gamma}^{g,h}$, and $\epsilon_j(g)$, etc., are defined as in the definition of the orbifold elliptic genus.

Our first goal is to prove

$$(3) \quad \mu_* E = \prod_{I_X} \frac{\frac{D_i}{2\pi i} \theta(\frac{D_i}{2\pi i} - (-\alpha_i + 1)z) \theta'(0)}{\theta(\frac{D_i}{2\pi i}) \theta(-(-\alpha_i + 1)z)}$$

To prove the above equality, we express both sides as the image under $\rho_{\hat{X}}$ and ρ_X of piece-wise convergent power series, and apply the push-forward formula from the previous section. To that end, let F be the piece-wise convergent power series defined as follows: Let $C_{\gamma}^{g,h}$ be the cone which corresponds to $X_{\gamma}^{g,h}$. For each cone $C = C_{J,j}$ containing $C_{\gamma}^{g,h}$, let $F_{\gamma}^{g,h}|_C =$

$$\frac{1}{|G|} \prod_J \frac{\frac{x_j^C}{2\pi i} \theta(\frac{x_j^C}{2\pi i} + \epsilon_j(g) - \epsilon_j(h)\tau - (-\beta_j + 1)z) \theta'(0)}{\theta(\frac{x_j^C}{2\pi i} + \epsilon_j(g) - \epsilon_j(h)\tau) \theta(-(-\beta_j + 1)z)} e^{2\pi i(-\beta_j + 1)\epsilon_j(h)z}.$$

For cones C not containing $C_\gamma^{g,h}$, we define $F_\gamma^{g,h}|_C = 0$. In the above expression, x_j^C are the piece-wise linear functions dual to the generators of C . If $\hat{D}_j$ are the divisors which correspond to the generators of C , then $\epsilon_j(g)$, etc., are the infinitesimal weights attached to the divisors, and β_j are the coefficients of $\hat{D}_j$. It is easy to see that $F_\gamma^{g,h}$ is a well-defined piece-wise convergent power series, and that $\rho_{\hat{X}}(F_\gamma^{g,h})$ is the $X_\gamma^{g,h}$ -th summand in the expression for E . We therefore define $F = \sum_{gh=hg, \gamma} F_\gamma^{g,h}$, so that $\rho_{\hat{X}}(F) = E$.

Similarly, define $H \in \mathbb{C}[[\Sigma_X]]$ to be the piece-wise convergent power series:

$$H|_C = \prod_{i=1}^{\dim C} \frac{\frac{x_i^C}{2\pi i} \theta(\frac{x_i^C}{2\pi i} - (-\alpha_i + 1)z) \theta'(0)}{\theta(\frac{x_i^C}{2\pi i}) \theta(-(-\alpha_i + 1)z)}.$$

Clearly $\rho_X(H)$ is equal to the right-hand side of 3. We are therefore reduced to proving that $\nu_* F = H$.

Let $C \subset \Sigma_X$ be a cone, and let Σ_α be the subdivisions of C lying in $\Sigma_{\hat{X}}$ which get mapped to C under ν . Let N_α denote the lattices of Σ_α , and N_C the lattice of Σ_C . Referring to the notation of section 5.3, the formula for ν_* tells us that:

$$(\nu_* F)_C = \sum_\alpha d_\alpha \sum_{C_j \subset \Sigma_\alpha} F|_{C_j} \frac{\prod_{i=1}^{\dim C} x_i^C}{\prod_{i=1}^{\dim C} x_i^{C_j}}.$$

For each $C_j \subset \Sigma_\alpha$ with the same dimension as C , note that $F_\gamma^{g,h}|_{C_j} \neq 0$ if and only if $C_\gamma^{g,h} \subset C_j$, i.e., if and only if g and h are elements of the group $G_\alpha = N_C/N_\alpha$. We may therefore write the push-forward of F as:

$$\begin{aligned} & \prod_{i=1}^{\dim C} x_i^C \sum_\alpha \frac{d_\alpha}{|G|} \sum_{g,h \in G_\alpha} \\ & \sum_{C_j \subset \Sigma_\alpha} \frac{\theta(\frac{x_i^{C_j}}{2\pi i} + \epsilon_i(g) - \epsilon_i(h)\tau - (-\beta_i + 1)z) \theta'(0)}{\theta(\frac{x_i^{C_j}}{2\pi i} + \epsilon_i(g) - \epsilon_i(h)\tau) \theta(-(-\beta_i + 1)z)} e^{2\pi i(-\beta_i + 1)z} \end{aligned}$$

By lemma 8.1 in [7], this is equal to $\sum_\alpha \frac{d_\alpha |G_\alpha|}{|G|} H|_C$. But since $\sum_\alpha d_\alpha |G_\alpha|$ describes the number of points in the pre-image of a generic point in a tubular neighborhood of the closed stratum corresponding to C , the coefficient

in front of $H|_C$ in the above expression is 1. This completes the proof of 3. To complete the proof, we apply the projection formula together with the following lemma relating the Chern classes of $\hat{X}$ to those of X :

Lemma 5.

$$\frac{c(T\hat{X})_T}{\mu^*c(TX)_T} = \frac{\prod_{I_{\hat{X}}} (1 + c_1(\hat{D}_j)_T)}{\prod_{I_X} (1 + c_1(D_i)_T)}$$

in $H_T^*(\hat{X})_{loc}$

For details, see [7]. The above lemma may be proved using an argument analogous to the proof of lemma 5 in [17]. $\square$

7. DEFORMATION TO THE NORMAL CONE

If Q is a holomorphic function in a neighborhood of the origin, then Q determines a map $\varphi_Q : K_T(\cdot) \rightarrow \hat{H}(\cdot)$ by the rule $E \mapsto \prod_i Q(e_i)$, where e_i represent the equivariant Chern roots of E . If, in addition, $Q(0) = 1$, then φ_Q is multiplicative in the sense that $\varphi_Q(E_1 \oplus E_2) = \varphi_Q(E_1)\varphi_Q(E_2)$.

More generally, let H be a finite abelian group with characters $\{\lambda\}$. Let $\{f_\lambda\}$ be an assignment of a holomorphic function in a neighborhood of the origin for each character. Such a collection $\{f_\lambda\}$ determines a map $\psi : K_T(\cdot) \otimes R(H) \rightarrow \hat{H}(\cdot)$ by the rule $\psi : E = \bigoplus_\lambda E_\lambda \mapsto \prod_\lambda \prod f_\lambda(e_{\lambda,i})$. Here $e_{\lambda,i}$ denote the equivariant Chern roots of the T -vectorbundle E_λ . Let us fix a multiplicative map $\varphi = \varphi_Q$ and a (possibly not multiplicative) map $\psi = \psi_{\{f_\lambda\}}$.

Let X be a compact $T \times H$ -variety and let Z be a smooth $T \times H$ -invariant subvariety. Let V be a connected component of X^H . In the proof of the lemma below, the only difficult case to examine is when $Z \cap V \equiv W$ is a proper subset of V . We therefore assume this throughout. Let $\pi : \tilde{X} \rightarrow X$ be the blow-up of X along Z and let $\tilde{V}$ be the proper transform of V . Clearly $N_{\tilde{V}/\tilde{X}}$ has the same H -character decomposition as $N_{V/X}$. We may therefore make sense of $\psi(N_{\tilde{V}/\tilde{X}})$. The goal of this section is to prove the following crucial technical lemma relating $\varphi(T\tilde{V})\psi(N_{\tilde{V}/\tilde{X}})$ to $\varphi(TV)\psi(N_{V/X})$.

Lemma 6.

$$\pi_*\varphi(T\tilde{V})\psi(N_{\tilde{V}/\tilde{X}}) - \varphi(TV)\psi(N_{V/X}) = (i_W)_*\Theta.$$

Here i_W is the inclusion map and $\Theta \in \hat{H}(W)$ is a universal class which depends only on the data of W , $N_{W/V}$, and $i_W^*N_{Z/X}$.

The argument given below is an adaptation of the argument in [16] which was given for the non-equivariant case with $\psi = 1$.

Proof. Let $\Pi : M_X \rightarrow X \times \mathbb{P}^1$ be the blow-up along $Z \times \{\infty\}$. We give M_X the obvious $T \times H$ action. Let M_V be the proper transform of $V \times \mathbb{P}^1$. Clearly $M_V \rightarrow V \times \mathbb{P}^1$ is the blow-up along $W \times \{\infty\}$. It is easy to see that $N_{M_V/M_X} \in K_T(M_V) \otimes R(H)$ has the same H -character decomposition as $N_{V/X}$.

Define N_0 to be the sub-bundle of $i_W^* N_{Z/X}$ on which H -acts trivially. Let $N_1 = i_W^* N_{Z/X} / N_0$. Then $i_W^* TX$ decomposes as $TW \oplus N_{W/Z} \oplus N_0 \oplus N_1$. Clearly $TW \oplus N_0 = i_W^* TV$, and therefore $N_{V/W} = N_0$ is a sub-bundle of $i_W^* N_{Z/X}$ with quotient N_1 .

Let $g : M_X \rightarrow \mathbb{P}^1$ be the composition $M_X \rightarrow X \times \mathbb{P}^1 \rightarrow \mathbb{P}^1$. Then $\text{div}(g) = X - \tilde{X} - \mathbb{P}(N_{Z/X} \oplus 1)$. Furthermore, $\text{div}(g|_{M_V}) = V - \tilde{V} - \mathbb{P}(N_{W/V} \oplus 1)$. Let i_V , $i_{\tilde{V}}$, and $i_{\mathbb{P}(N_{W/V} \oplus 1)}$ denote the respective inclusion maps of these divisors in M_V .

Let $p : \mathbb{P}(N_{W/V} \oplus 1) \rightarrow W$ be the obvious fibration, and let S denote the tautological bundle over $\mathbb{P}(N_{W/V} \oplus 1)$.

CLAIM:

$$\begin{aligned} (4) \quad & i_V^* N_{M_V/M_X} = N_{V/X} \\ (5) \quad & i_{\tilde{V}}^* N_{M_V/M_X} = N_{\tilde{V}/\tilde{X}} \\ (6) \quad & i_{\mathbb{P}(N_{W/V} \oplus 1)}^* N_{M_V/M_X} = p^* N_{W/Z} \oplus p^* N_1 \otimes S^* \end{aligned}$$

(1) is obvious. To prove (2), note first that $TM_X|_{\tilde{V}}$ decomposes as $T\tilde{V} \oplus N_{\tilde{V}/\tilde{X}} \oplus N_{\tilde{X}/M_X}|_{\tilde{V}}$ and also as $T\tilde{V} \oplus N_{\tilde{V}/M_V} \oplus N_{M_V/M_X}|_{\tilde{V}}$. Then simply notice that in both decompositions, $N_{\tilde{V}/\tilde{X}}$ and $i_{\tilde{V}}^* N_{M_V/M_X}$ are the nontrivial H -eigenspaces.

To prove (3), note that $TM_X|_{\mathbb{P}(N_{W/V} \oplus 1)}$ decomposes in the following two ways:

$$\begin{aligned} TM_X|_{\mathbb{P}(N_{W/V} \oplus 1)} &= \\ T\mathbb{P}(N_{W/V} \oplus 1) \oplus \mathcal{O}(-1)_{\mathbb{P}(N_{W/V} \oplus 1)} \oplus i_{\mathbb{P}(N_{W/V} \oplus 1)}^* N_{M_V/M_X} &= \\ T\mathbb{P}(N_{W/V} \oplus 1) \oplus N_{\mathbb{P}(N_{W/V} \oplus 1)/\mathbb{P}(N_{Z/X} \oplus 1)} \oplus i_{\mathbb{P}(N_{W/V} \oplus 1)}^* \mathcal{O}(-1)_{\mathbb{P}(N_{Z/X} \oplus 1)} \end{aligned}$$

Observing $i_{\mathbb{P}(N_{W/V} \oplus 1)}^* \mathcal{O}(-1)_{\mathbb{P}(N_{Z/X} \oplus 1)} = \mathcal{O}(-1)_{\mathbb{P}(N_{W/V} \oplus 1)}$ it follows that $i_{\mathbb{P}(N_{W/V} \oplus 1)}^* N_{M_V/M_X} = N_{\mathbb{P}(N_{W/V} \oplus 1)/\mathbb{P}(N_{Z/X} \oplus 1)}$. It is easy to verify that this bundle in turn is equal to $p^* N_{W/Z} \oplus p^* N_1 \otimes S^*$.

Since $\text{div}(g|_{M_V}) = V - \tilde{V} - \mathbb{P}(N_{W/V} \oplus 1)$, as equivariant classes $V = \tilde{V} + \mathbb{P}(N_{W/V} \oplus 1)$. Let u be the equivariant Thom class of $\tilde{V}$ (that is, the Thom class of its normal bundle), and let v be the equivariant Thom class of $\mathbb{P}(N_{W/V} \oplus 1)$. Then $u + v$ is the equivariant Thom class of V . Since

V is disjoint from $\tilde{V}$ and $\mathbb{P}(N_{W/V} \oplus 1)$, we have the relations $u(u+v) = v(u+v) = 0$. Note also that uv is the equivariant Thom class of $\mathbb{P}(N_{W/V})$, which is the exceptional divisor of $\tilde{V} \rightarrow V$.

Let f be the holomorphic function in a neighborhood of the origin which satisfies the relation $Q(z) = \frac{z}{f(z)}$. Then by the above claim:

$$\begin{aligned}\varphi(TM_V)\psi(N_{M_V/M_X})f(u+v) &= (i_V)_*\varphi(TV)\psi(N_{V/X}) \\ \varphi(TM_V)\psi(N_{M_V/M_X})f(u) &= (i_{\tilde{V}})_*\varphi(T\tilde{V})\psi(N_{\tilde{V}/\tilde{X}}) \\ \varphi(TM_V)f(v) &= (i_{\mathbb{P}(N_{W/V} \oplus 1)})_*\varphi(T\mathbb{P}(N_{W/V} \oplus 1))\psi(N_{\mathbb{P}(N_{W/V} \oplus 1)/\mathbb{P}(N_{Z/X} \oplus 1)})\end{aligned}$$

Note that since u and v are equivariant Thom classes, their degree zero part is an element of $C^\infty(M_V) \otimes \mathfrak{t}^*$ which vanishes at the origin of $\mathfrak{t}$. Hence $f(u)$, etc., are well-defined elements of $\hat{H}(M_V)$.

Since $f(z) = z + \dots$, we can define $g = f^{-1}$ in a possibly smaller neighborhood of the origin. Consider the two-variable holomorphic function $h(z_1, z_2) = f(z_1 + z_2)$. Clearly $h(z_1, z_2) = h(g(f(z_1)), g(f(z_2)))$. Define $F(y_1, y_2) = h(g(y_1), g(y_2))$. Then F is holomorphic in a neighborhood of the origin and $F(f(z_1), f(z_2)) = f(z_1 + z_2)$. From [11] we have the formula:

$$F(y_1, y_2)g'(y_1)g'(y_2) = \sum_{(i,j) \neq (0,0)} \varphi(H_{ij})y_1^i y_2^j.$$

Here $\varphi(H_{ij})$ is the non-equivariant genus induced by f of the degree $(1,1)$ hypersurface $H_{ij} \subset \mathbb{P}^i \times \mathbb{P}^j$. If we plug in $f(u)$ and $f(v)$ for y_1 and y_2 , we get $F(f(u), f(v))g'(f(u))g'(f(v)) = f(u+v)g'(f(u))g'(f(v))$. By the relations $u(u+v) = v(u+v) = 0$, this last term is cohomologous to $f(u+v)$. It is instructive to go over this last point in detail. Write $f(x+y) = \sum a_n(x+y)^n$ and $g'(f(x))g'(f(y)) = \sum b_{ij}x^i y^j$. Note that $b_{00} = 1$. Let $h_{NIJ} = \sum_{n \leq N} a_n(x+y)^n \cdot \sum_{i \leq I, j \leq J} b_{ij}x^i y^j$. Then $h_{NIJ} \rightarrow f(x+y)g'(f(x))g'(f(y))$ in the C^∞ topology. Abusing notation, let u and v denote fixed representatives for their respective cohomology classes. We have that $h_{NIJ}(u, v) = \sum_{n \leq N} a_n(u+v)^n + d\eta_{NIJ}$. Evaluating at a sufficiently small s gives: $h_{NIJ}(u, v)(s) = \sum_{n \leq N} a_n(u(s) + v(s))^n + d\eta_{NIJ}(s)$. Taking the limit in the indices N, I, J , we get $h(u, v)(s) = f(u(s) + v(s)) + d\eta(s)$. We therefore have that $ev(h) = ev(f(u+v))$, and therefore $h = f(u+v)$ in $\hat{H}(M_V)$.

At the same time, $f(u+v) = \sum_{(i,j) \neq (0,0)} \varphi(H_{ij})f(u)^i f(v)^j$. Thus

$$\begin{aligned} \varphi(TM_V)\psi(N_{M_V/M_X})f(u+v) &= \varphi(TM_V)\psi(N_{M_V/M_X})f(u) + \\ \varphi(TM_V)\psi(N_{M_V/M_X})f(v) &+ \varphi(TM_V)\psi(N_{M_V/M_X}) \sum_{i+j \geq 2} \varphi(H_{ij})f(u)^i f(v)^j \end{aligned}$$

From the relations $u^2 = -uv$ and $v^2 = -uv$, we have that $f(u)^{i+1} = f(u)f(-v)^i$. Therefore,

$$\begin{aligned} \sum_{i+j \geq 2} \varphi(H_{ij})f(u)^i f(v)^j &= \\ \sum_{i+j \geq 2, i \geq 1} \varphi(H_{ij})f(u)f(-v)^{i-1} f(v)^j &+ \sum_{j \geq 2} \varphi(H_{0j})f(v)^j \end{aligned}$$

We therefore have $\sum_{i+j \geq 2} \varphi(H_{ij})f(u)^i f(v)^j = uv \frac{f(u)}{u} G(v) + vJ(v)$ for some universal convergent power series G and J . Let $\nu = i_{\mathbb{P}(N_{W/V} \oplus 1)}^* v$. Clearly $\nu = c_1(S)$. Let $w = \nu|_{\mathbb{P}(N_{W/V})}$. Finally, for ease of notation, write $N = p^* N_{W/Z} \oplus p^* N_1 \otimes S^*$.

$$\begin{aligned} \varphi(TM_V)\psi(N_{M_V/M_X})f(u+v) &= \\ \varphi(TM_V)\psi(N_{M_V/M_X})f(u) &+ \varphi(TM_V)\psi(N_{M_V/M_X})f(v) + \\ \varphi(TM_V)\psi(N_{M_V/M_X})uv \frac{f(u)}{u} G(v) &+ \varphi(TM_V)\psi(N_{M_V/M_X})vJ(v) \end{aligned}$$

It follows that

$$\begin{aligned} (i_V)_* \varphi(TV)\psi(N_{V/X}) &= \\ (i_{\tilde{V}})_* \varphi(T\tilde{V})\psi(N_{\tilde{V}/\tilde{X}}) &+ (i_{\mathbb{P}(N_{W/V} \oplus 1)})_* \varphi(T\mathbb{P}(N_{W/V} \oplus 1))\psi(N) + \\ (i_{\mathbb{P}(N_{W/V})})_* \varphi(T\mathbb{P}(N_{W/V}) \oplus S|_{\mathbb{P}(N_{W/V})}) &\psi(N|_{\mathbb{P}(N_{W/V})})G(w) + \\ (i_{\mathbb{P}(N_{W/V} \oplus 1)})_* \varphi(T\mathbb{P}(N_{W/V} \oplus 1) \oplus S) &\psi(N)J(\nu) \end{aligned}$$

Now apply the push-forward Π_* to the above equation. Note that $\Pi \circ i_V$ is the inclusion $v \mapsto (v, 0)$ in $V \times \mathbb{P}^1$. $\Pi \circ i_{\tilde{V}}$ is the composition of the blow-down $\tilde{V} \rightarrow V$ with the inclusion of V at infinity in $V \times \mathbb{P}^1$. From the blow-up diagram:

$$\begin{array}{ccc}
\mathbb{P}(N_{W/V} \oplus 1) & \xrightarrow{i_{\mathbb{P}(N_{W/V} \oplus 1)}} & M_V \\
\hat{\Pi} \downarrow & & \downarrow \Pi \\
W \times \{\infty\} & \xrightarrow{i_W} & V \times \mathbb{P}^1
\end{array}$$

we have that $\Pi \circ i_{\mathbb{P}(N_{W/V} \oplus 1)} = i_W \circ \hat{\Pi}$. Finally, $\Pi \circ i_{\mathbb{P}(N_{W/V})}$ is clearly the composition of the blow-down map $\hat{\Pi} : \mathbb{P}(N_{W/V}) \rightarrow W$ with the inclusion i_W . Therefore, applying the pushforward Π_* gives the equation:

$$\begin{aligned}
& \varphi(TV)\psi(N_{V/X}) = \\
& \pi_*\varphi(T\tilde{V})\psi(N_{\tilde{V}/\tilde{X}}) + (i_W)_* \left\{ \hat{\Pi}_*\varphi(T\mathbb{P}(N_{W/V} \oplus 1))\psi(N) + \right. \\
& \hat{\pi}_*\varphi(T\mathbb{P}(N_{W/V}) \oplus S|_{\mathbb{P}(N_{W/V})})\psi(N|_{\mathbb{P}(N_{W/V})})G(w) + \\
& \left. \hat{\Pi}_*\varphi(T\mathbb{P}(N_{W/V} \oplus 1) \oplus S)\psi(N)J(\nu) \right\}
\end{aligned}$$

Since the term in the curly braces depends only on the data of $i_W^*N_{Z/X}$, $N_{W/V}$, and W , this proves the lemma. $\square$

Remark 3. Of course in the above proof, if $Q(x) = \frac{\frac{x}{2\pi i}\theta(\frac{x}{2\pi i}-z)\frac{\theta'(0)}{2\pi i}}{\theta(\frac{x}{2\pi i})\theta(-z)}$ and $f_\lambda = \frac{e^{2\pi i\lambda(h)\tau z}\theta(\frac{x}{2\pi i}+\lambda(g)-\lambda(h)\tau-z)\frac{\theta'(0)}{2\pi i}}{\theta(\frac{x}{2\pi i}+\lambda(g)-\lambda(h)\tau)\theta(-z)}$, then $\varphi(TV)\psi(N_{V/X})$ is the equivariant elliptic class associated to the pair (g, h) and the (g, h) -fixed component V .

8. THE NORMAL CONE SPACE

Let $W \subset X$ be a connected T -invariant subvariety of a projective T -space X . Suppose the normal bundle $N_{W/X}$ splits into a composition $L_1 \oplus \dots \oplus L_k$ of T -vectorbundles. Define $p : X^* \rightarrow W$ to be the fiber bundle with fiber $p^{-1}(w) = \mathbb{P}(L_1 \oplus 1)_w \times \dots \times \mathbb{P}(L_k \oplus 1)_w$. It is easy to see that X^* contains a copy of W with the same normal bundle $N_{W/X}$. In our proof of the blow-up formula, we will ultimately reduce all computations on X to computations on the more manageable space X^* . We therefore devote this section to gathering some important facts about the topology of X^* .

If we give the trivial vectorbundle 1 the trivial action, then the action of T on W lifts naturally to X^* . Give L_i a metric so that T acts on L_i by isometries, and give $\mathbb{P}(L_i \oplus 1)$ the induced metric.

Define vectorbundles $Q_i \rightarrow X^*$ as follows: For $w \in W$ and $(\ell_1, \dots, \ell_k)$ lines in $(L_1 \oplus 1)_w, \dots, (L_k \oplus 1)_w$, define $(Q_i)_{(w, \ell_1, \dots, \ell_k)} = \ell_i^\perp \subset (L_i \oplus 1)_w$.

These bundles inherit natural T -actions. Observe furthermore that $i_W^*(Q_i)$ is naturally isomorphic to L_i .

Define $V_i \subset X^*$ to be the subvariety $\{(w, \ell_1, \dots, \ell_k) : \ell_i = [0 : 1]\}$. For $i = 1, \dots, k$, V_i are connected T -subvarieties, with connected intersection locus, and $W = V_1 \cap \dots \cap V_k$.

Finally, let $f : \widetilde{X^*} \rightarrow X^*$ denote the blow-up of X^* along $Z = V_1 \cap \dots \cap V_j$ with exceptional divisor E . We have the following intersection-theoretic result:

Theorem 7. $c_{top}(f^*Q_1 \oplus \dots \oplus f^*Q_j \otimes \mathcal{O}(-E))_T = 0$.

Proof. We will show that $\text{Hom}(L, f^*Q_1 \oplus \dots \oplus f^*Q_j)$ has an equivariant global nowhere zero section, where L is a line bundle with the same equivariant first Chern class as $\mathcal{O}(E)$.

We first give an explicit construction of the line bundle L . Let $0 \leq t_1, \dots, t_j$. Define $S_{t_1, \dots, t_j} \subset \widetilde{X^*}$ to be the subset $\{(w, [v_1 : 1], \dots, [v_j : 1], [v_1 : \dots : v_j], \bar{\ell}) : \|v_i\| = t_i\}$. Here $\bar{\ell}$ represents a point in $\mathbb{P}(L_{j+1} \oplus 1)_w \times \dots \times \mathbb{P}(L_k \oplus 1)_w$. We will refer to points in $\widetilde{X^*}$ which are not contained in any $S_{t_1, \dots, t_j}$ as *points at infinity*. Let $\rho : [0, \infty) \rightarrow \mathbb{R}$ be a bump function equal to zero in the region $[0, 1/3)$ and equal to one in the region $[2/3, \infty)$. For $0 \leq t_i \leq 1$ and $v = (w, [v_1 : 1], \dots, [v_j : 1], [v_1 : \dots : v_j], \bar{\ell})$ a point in $S_{t_1, \dots, t_j}$, let $L_v \subset (L_1 \oplus 1)_w \oplus \dots \oplus (L_j \oplus 1)_w$ be the span of the vector $\tilde{v} = ((1 - \rho(t_1))v_1, \rho(t_1), \dots, (1 - \rho(t_j))v_j, \rho(t_j))$. Outside this set, we define L_v to be the span of the vector $0 \oplus 1 \oplus \dots \oplus 0 \oplus 1$. This clearly defines a smooth line bundle on $\widetilde{X^*}$ with the same equivariant first Chern class as $\mathcal{O}(E)$.

We now prove that $\text{Hom}(L, f^*Q_1 \oplus \dots \oplus f^*Q_j)$ has a global equivariant nowhere zero section. For $v_i \in L_i$, define:

$$h_i(v_i) = \begin{cases} (-v_i, \|v_i\|^2) & \|v_i\| \leq 1 \\ (-\frac{v_i}{\|v_i\|^2}, 1) & \|v_i\| \geq 1 \end{cases}$$

For $v = (w, [v_1 : 1], \dots, [v_j : 1], [v_1 : \dots : v_j], \bar{\ell}) \in S_{t_1, \dots, t_j}$, we define $L_v \rightarrow (Q_1 \oplus \dots \oplus Q_j)_{f(v)}$ by $\tilde{v} \mapsto (h_1(v_1), \dots, h_j(v_j))$. This section extends in a natural way to the points at infinity, giving us a *continuous* nowhere zero equivariant section s_0 of $\text{Hom}(L, f^*Q_1 \oplus \dots \oplus f^*Q_j)$. The section is only continuous because the function:

$$h(x) = \begin{cases} x^2 & |x| \leq 1 \\ \frac{1}{x^2} & |x| \geq 1 \end{cases}$$

is not smooth. However, we may remedy this by approximating our continuous section s_0 by a smooth section and then averaging over the group T .

Since s_0 was nowhere zero and fixed by the T -averaging process, the new smooth section will remain nowhere zero after averaging over T . $\square$

Remark 4. The intuition behind the preceding theorem is that if $Z = D_1 \cap \dots \cap D_j$ is the complete intersection of a collection of normal crossing divisors, then the proper transforms $\tilde{D}_i$ of D_i are disjoint when we blow up along Z . We therefore have that $c_{top}(\mathcal{O}(\tilde{D}_1) \oplus \dots \oplus \mathcal{O}(\tilde{D}_j))_T = 0$. While the above theorem is known, the proof given here will be useful later.

One easily observes that $i_{V_i}^* Q_i = N_{V_i/X^*}$. We might expect, therefore, that $c_{top}(Q_i)_T$ is the equivariant Thom class of V_i . This is in fact the case, as the following lemma proves:

Lemma 7. $c_{top}(Q_i)_T = i_{V_i*} 1$

Proof. By the naturality properties of the equivariant Chern classes, it is enough to prove the above lemma for $X^* = \mathbb{P}(L \oplus 1)$, where L is a T -vectorbundle over W , and Q is the universal quotient bundle over $\mathbb{P}(L \oplus 1)$. Here, W itself plays the role of V_i in the statement of the lemma. As above, we endow the trivial bundle 1 with the trivial T -action.

Let us first prove that $c_{top}(Q_i) = i_{W*} 1$ in the non-equivariant category. Let $p : \mathbb{P}(L \oplus 1) \rightarrow W$ denote the obvious projection map. Let $r = \text{rk}(Q)$. From the exact sequence:

$$0 \rightarrow S \rightarrow p^*(L \oplus 1) \rightarrow Q \rightarrow 0$$

we have that $c_r(Q) = \left(\frac{p^*c(L)}{1 - c_1(S^*)} \right)_r$, where $(\cdot)_r$ denotes the degree r part. Let $p_w : \mathbb{P}(L \oplus 1)_w \rightarrow w$ denote the restriction of p to the fiber over w . Then $c_r(Q)|_{\mathbb{P}(L \oplus 1)_w} = c_1(S^*)^r|_{\mathbb{P}(L \oplus 1)_w}$, which clearly integrates to 1 over the fiber.

We next observe that Q has a global no-where zero section away from W . We define this section as follows: $s : (w, [v : z]) \mapsto (w, [v : z], \frac{\bar{z}v}{\|v\|^2}, -1)$. We therefore have that $c_r(Q)$ is exact away from W . Hence, by subtracting off an exact form, we may represent $c_r(Q)$ by a form which has compact support in a tubular neighborhood of W , and which integrates to one along every fiber of the tubular neighborhood. It follows that $c_r(Q) = i_{W*} 1$ at least in the non-equivariant sense.

In general, $c_r(Q)_T$ is at least an equivariant extension of the Thom class of W . Moreover, $c_r(Q)_T$ is equivariantly exact outside W . We prove this as follows: Observe first that the non-zero section we defined on the complement of W was in fact equivariant. This section therefore induces a splitting $Q = Q' \oplus \mathbb{C}$ outside of W . Since we endowed 1 with the trivial T -action, one may easily verify that the $\mathbb{C}$ in the above splitting inherits a trivial action as well. It follows that $c_r(Q)_T$ is equivariantly exact outside of W .

By subtracting off an exact form, we get that $c_r(Q)_T$ may be represented by an equivariant form which has compact support in a tubular neighborhood of W , and which integrates to one along every fiber of the tubular neighborhood. It follows that $c_r(Q)_T$ is the equivariant Thom class of W . $\square$

We next prove a formula relating the equivariant Chern class of the blow-up of X^* along Z to that of X^* . Note first that TX^* splits holomorphically and equivariantly into a direct sum of sub-bundles $F \oplus M$, with $i_Z^*F = TZ$ and $i_Z^*M = N_{Z/X^*}$. We may therefore apply the following lemma to compare the equivariant Chern classes of X^* and $\text{Bl}_Z X^*$.

Lemma 8. *Let Y be a complex T -space, and $Z \subset Y$ a T -invariant complex submanifold. Suppose that TY splits holomorphically and equivariantly into a direct sum of sub-bundles $F \oplus M$ such that $i_Z^*F = TZ$ and $i_Z^*M = N_{Z/Y}$. Let $f : \tilde{Y} \rightarrow Y$ be the blow-up of Y along Z with exceptional divisor E . Then:*

$$(7) \quad c(T\tilde{Y})_T = c(f^*F)_T c(f^*M \otimes \mathcal{O}(-E))_T c(\mathcal{O}(E))_T$$

in the ring $H_T^*(\tilde{Y})_{loc}$.

Proof. By localization, it suffices to prove the equality at every fixed component in $\tilde{Y}$. Let $\tilde{P} \subset \tilde{Y}$ be a fixed component which is the proper transform of a fixed component $P \subset Y$. If P is disjoint from Z , then the equality of (4) at $\tilde{P} \cong P$ is trivial. Otherwise, $\tilde{P}$ is equal to the blow-up of P at $P \cap Z$. Note that i_P^*F decomposes as $F_0 \oplus F_1$, where T acts trivially on F_0 and nontrivially on F_1 . Similarly, $i_P^*M = M_0 \oplus M_1$. Clearly $TP = F_0 \oplus M_0$ and $N_{P/Y} = F_1 \oplus M_1$.

Applying $i_{\tilde{P}}^*$ to (4), we have:

$$\begin{aligned} i_{\tilde{P}}^*(\text{LHS}) &= c(T\tilde{P})c(N_{\tilde{P}/\tilde{Y}})_T \\ i_{\tilde{P}}^*(\text{RHS}) &= c(f^*F_0)c(f^*M_0 \otimes \mathcal{O}(-E))c(i_{\tilde{P}}^*\mathcal{O}(E)) \\ &\quad c(f^*F_1)_T c(f^*M_1 \otimes \mathcal{O}(-E))_T \end{aligned}$$

Since $\tilde{P}$ is the blow-up of P along $P \cap Z$ and $i_{P \cap Z}^*M_0 = N_{P \cap Z/P}$, the relation $c(T\tilde{P}) = c(f^*F_0)c(f^*M_0 \otimes \mathcal{O}(-E))c(i_{\tilde{P}}^*\mathcal{O}(E))$ is well-known (see [9]). It suffices therefore to prove that $c(N_{\tilde{P}/\tilde{Y}})_T = i_{\tilde{P}}^*(c(f^*F_1)_T c(f^*M_1 \otimes \mathcal{O}(-E))_T)$. To this end we prove the following claim:

CLAIM: $N_{\tilde{P}/\tilde{Y}} \cong i_{\tilde{P}}^*(f^*F_1 \oplus f^*M_1 \otimes \mathcal{O}(-E))$ as T -vectorbundles.

To prove this, consider f as a map $f : \tilde{P} \rightarrow P$. For simplicity of notation, write E for $E \cap \tilde{P}$. View $N_{\tilde{P}/\tilde{Y}}$ as a sheaf, i.e., $N_{\tilde{P}/\tilde{Y}}(U) = \Gamma(U, N_{\tilde{P}/\tilde{Y}})$. The derivative $Df : N_{\tilde{P}/\tilde{Y}} \rightarrow f^*N_{P/Y} = f^*F_1 \oplus f^*M_1$ is a sheaf map, and maps onto the subsheaf $f^*F_1 \oplus f^*M_1(-E)$. Here $f^*M_1(-E)$ represents the subsheaf of f^*M_1 corresponding to sections of f^*M_1 which vanish along E . Let s_0 denote the global section of $\mathcal{O}(E)$ induced by the defining equations of E . Then $\otimes s_0^{-1} : f^*M_1(-E) \rightarrow f^*M_1 \otimes \mathcal{O}(-E)$ is a sheaf isomorphism. Define $\beta = (id \oplus \otimes s_0^{-1}) \circ Df$. By computing in local coordinates, one verifies easily that $\beta : N_{\tilde{P}/\tilde{Y}} \rightarrow f^*F_1 \oplus f^*M_1 \otimes \mathcal{O}(-E)$ is a sheaf isomorphism. Furthermore, for s a local section of $N_{\tilde{P}/\tilde{Y}}$, $s(p) = 0$ if and only if $(\beta s)(p) = 0$. Therefore, β induces an isomorphism of the corresponding vectorbundles. Moreover, since β is clearly equivariant, the vectorbundle isomorphism is equivariant.

This completes the proof if $P \cap E < P$. Next, suppose that $\tilde{P} \subset E$. For this case, it suffices to prove the following fact:

CLAIM $i_E^*(T\tilde{Y} \oplus \mathbb{C}) = i_E^*(f^*F \oplus f^*M \otimes \mathcal{O}(-E) \oplus \mathcal{O}(E))$.

We prove this as follows: Let $\pi : \mathbb{P}(N_{Z/Y}) \rightarrow Z$ be the natural projection, and let S denote the tautological bundle over $\mathbb{P}(N_{Z/Y})$. Then $i_E^*T\tilde{Y} = T\mathbb{P}(N_{Z/Y}) \oplus S$.

Since $i_E^*f^* = \pi^*i_Z^*$, we have that $i_E^*f^*F = \pi^*i_Z^*F = \pi^*TZ$ and $i_E^*f^*M = \pi^*N_{Z/Y}$. Thus,

$$i_E^*(f^*F \oplus f^*M \otimes \mathcal{O}(-E) \oplus \mathcal{O}(E)) = \pi^*TZ \oplus \pi^*N_{Z/Y} \otimes S^* \oplus S$$

From the exact sequence $0 \rightarrow S \rightarrow \pi^*N_{Z/Y} \rightarrow Q \rightarrow 0$, we have that $\pi^*N_{Z/Y} \otimes S^* = \mathbb{C} \oplus Q \otimes S^*$, where Q is the tautological quotient bundle. The claim then follows from the observation that $T\mathbb{P}(N_{Z/Y}) = \pi^*TZ \oplus Q \otimes S^*$. This completes the proof of the lemma. $\square$

Remark 5. Note that if $\tilde{Y}$ is equivariantly formal, the above proof implies that (4) holds in the unlocalized ring $H_T^*(\tilde{Y})$.

We may rewrite the left-hand side of (4) as $c(f^*TY)_T c(f^*M)_T^{-1} c(f^*M \otimes \mathcal{O}(-E))_T c(\mathcal{O}(E))_T$. Viewed as an element of $\hat{H}(\tilde{Y})$, it is easy to verify that this expression remains the same if we replace M by any bundle M' with $i_Z^*M' = N_{Z/Y}$. We therefore obtain the following corollary pertaining to X^* :

Corollary 1. *Let $f : \tilde{X}^* \rightarrow X^*$ be the blow-up of X^* along $Z = V_1 \cap \dots \cap V_j$ for $j \leq k$ with exceptional divisor E . Then the following formula holds in $\hat{H}(\tilde{X}^*)$:*

$$c(T\widetilde{X}^*)_T = \frac{c(f^*TX^*)_T}{\prod_{i=1}^j c(f^*Q_i)_T} \prod_{i=1}^j c(f^*Q_i \otimes \mathcal{O}(-E))_T c(\mathcal{O}(E))_T$$

We end this section with a technical lemma which is the blow-up analogue of lemma 7.

Lemma 9. *For $1 \leq i \leq k$, let $\widetilde{V}_i$ be the proper transform of V_i under the above blow-up: $f: \widetilde{X}^* \rightarrow X^*$. Then $c_{top}(f^*Q_i \otimes \mathcal{O}(-E))_T$ is the equivariant Thom class of $\widetilde{V}_i$ for $1 \leq i \leq j$, and $c_{top}(f^*Q_i)_T$ is the equivariant Thom class of $\widetilde{V}_i$ for $j+1 \leq i \leq k$. Moreover $i_{\widetilde{V}_i}^* c(f^*Q_i \otimes \mathcal{O}(-E))_T = c(N_{\widetilde{V}_i/\widetilde{X}^*})_T$ for $1 \leq i \leq j$ and $i_{\widetilde{V}_i}^* c(f^*Q_i)_T = c(N_{\widetilde{V}_i/\widetilde{X}^*})_T$ for $j+1 \leq i \leq k$.*

Proof. Let $1 \leq i \leq j$. Recall the equivariant continuous nowhere vanishing section s_0 of $\text{Hom}(L, f^*Q_1 \oplus \dots \oplus f^*Q_k)$ constructed in the proof of theorem 8. Let π_i denote the projection from $f^*Q_1 \oplus \dots \oplus f^*Q_k$ onto the i -th factor. Then $\pi_i \circ s_0$ is an equivariant section of $f^*Q_i \otimes L^*$. From the construction of s_0 , it is clear that s_0 vanishes precisely along $\widetilde{V}_i$. Hence, the equivariant top Chern class of $f^*Q_i \otimes L^*$ is the equivariant Thom class of $\widetilde{V}_i$. Since L has the same equivariant first Chern class as $\mathcal{O}(E)$, this proves $c_{top}(f^*Q_i \otimes \mathcal{O}(-E))_T = i_{\widetilde{V}_i}^* 1$.

To prove the second part of the lemma for $1 \leq i \leq j$, apply $i_{\widetilde{V}_i}^*$ to both sides of the equation in corollary 1. The LHS becomes $c(T\widetilde{V}_i)_T c(N_{\widetilde{V}_i/\widetilde{X}^*})_T$ while the RHS becomes:

$$\frac{c(f^*N_{V_i/X^*} \otimes \mathcal{O}(-E \cap \widetilde{V}_i))_T c(f^*TV_i)_T}{\prod_{\ell \neq i}^j c(f^*N_{V_\ell/X^*})_T} \times \prod_{\ell \neq i}^j c(f^*N_{V_\ell/X^*} \otimes \mathcal{O}(-E \cap \widetilde{V}_i))_T c(\mathcal{O}(E \cap \widetilde{V}_i))_T$$

Here we have used the fact that $i_{V_\ell}^* Q_\ell = N_{V_\ell/X^*}$. By corollary 1, the factor multiplying $i_{\widetilde{V}_i}^* c(f^*Q_i \otimes \mathcal{O}(-E))_T = c(f^*N_{V_i/X^*} \otimes \mathcal{O}(-E \cap \widetilde{V}_i))_T$ in the above expression is equal to $c(T\widetilde{V}_i)_T$. We therefore have that $i_{\widetilde{V}_i}^* c(f^*Q_i \otimes \mathcal{O}(-E))_T = c(N_{\widetilde{V}_i/\widetilde{X}^*})_T$.

Next, let $j+1 \leq i \leq k$. From the proof of lemma 7 we know that Q_i has an equivariant nowhere zero section in the complement of V_i . Pulling back this section by f gives an equivariant nowhere zero section in the complement of $f^{-1}V_i = \widetilde{V}_i$. It follows that $c_{top}(f^*Q_i)_T$ must be localized in

a neighborhood of $\widetilde{V}_i$. By the equivariant Thom isomorphism, $c_{top}(f^*Q_i)_T$ must be a complex number multiple of the equivariant Thom class of $\widetilde{V}_i$. But since $f_*c_{top}(f^*Q_i)_T$ is the Thom class of V_i , this complex number multiple must be equal to one.

The proof of the second part of the lemma for $j+1 \leq i \leq k$ is analogous to the proof given above for $1 \leq i \leq j$. $\square$

9. TWISTED POLYHEDRAL COMPLEX

Throughout, we assume the following:

- (a) X is a smooth projective variety with a holomorphic $T \times H$ action, where T is a torus and H is a finite abelian group.
- (b) $Z \subset X$ is a T -invariant smooth subvariety.
- (c) $V \subset X^H$ is a connected component.
- (d) $W = V \cap Z$ is connected. (see remark at the end of this section)

As noted above, $i_W^*N_{Z/X}$ splits as $N_0 \oplus N_1$, where N_0 denotes the sub-bundle of $i_W^*N_{Z/X}$ on which H acts trivially. Furthermore, N_1 decomposes as a sum of sub-bundles $\oplus N_\lambda$ corresponding to the characters of the H -action on N_1 . Finally, $N_{W/Z}$ also splits into character sub-bundles $\oplus N_\varepsilon$.

Let $E_1, \dots, E_\ell, D_1, \dots, D_k$ be smooth normal crossing divisors on X intersecting Z normally and non-trivially. We label these divisors so that $i_W^*\mathcal{L}_{D_i} \hookrightarrow N_0$ and $i_W^*\mathcal{L}_{E_j} \hookrightarrow N_1$. Write $i_W^*\mathcal{L}_{D_i} = \Delta_i$ and $i_W^*\mathcal{L}_{E_j} = \xi_j$. Write $N_0 = F_0 \oplus \bigoplus \Delta_i$ and $N_1 = \bigoplus F_\lambda \oplus \bigoplus \xi_j$.

Define a new space X^* as follows: $p : X^* \rightarrow W$ is the fiber bundle with fiber $p^{-1}(w) =$

$$\prod \mathbb{P}(N_\varepsilon \oplus 1)_w \times \mathbb{P}(F_0 \oplus 1)_w \times \prod \mathbb{P}(\Delta_i \oplus 1)_w \times \prod \mathbb{P}(F_\lambda \oplus 1)_w \times \prod \mathbb{P}(\xi_j \oplus 1)_w$$

Clearly X^* contains a copy of W with normal bundle $N_{W/X}$. Moreover, as in the previous section, each of the bundles $N_\varepsilon, F_0, F_\lambda, \Delta_i$, and ξ_j are the restrictions to W of global equivariant bundles $Q_{N_\varepsilon}, Q_{F_0}, Q_{F_\lambda}, Q_{\Delta_i}$ and Q_{ξ_j} . Recall also that the top Chern classes of these bundles are Poincaré dual to varieties $V_{N_\varepsilon}, V_{F_0}, V_{F_\lambda}, V_{\Delta_i}$, and V_{ξ_j} . The analogue of Z in X^* is the intersection locus of the varieties $V_{F_0}, V_{F_\lambda}, V_{\Delta_i}$, and V_{ξ_j} . We will continue to call this intersection locus Z .

We associate a polyhedral complex with integral structure (Σ, N_Σ) to X^* as follows: Let $s_\varepsilon = \text{rk}(N_\varepsilon)$, $r = \text{rk}(F_0)$, and let $r_\lambda = \text{rk}(F_\lambda)$. Let $w^{\varepsilon, a}, x^b, y^{\lambda, c}, d^i$, and e^j be an integral basis for $N_\Sigma = \mathbb{Z}^{\sum_\varepsilon s_\varepsilon + r + \sum_\lambda r_\lambda + k + \ell}$, for $a = 1, \dots, s_\varepsilon$, $b = 1, \dots, r$, $c = 1, \dots, r_\lambda$, $i = 1, \dots, k$, and $j = 1, \dots, \ell$. Define Σ to be the cone in the first orthant of the vectorspace $N_\Sigma \otimes \mathbb{R}$.

Let $w_{\varepsilon,a}, x_b, y_{\lambda,c}, d_i$, and e_j be the linear forms on this vectorspace which are dual to above basis vectors. Then $\mathbb{C}[\Sigma] = \mathbb{C}[w_{\varepsilon,a}, x_b, y_{\lambda,c}, d_i, e_j]$. Let $G = \prod_{\varepsilon} S_{s_{\varepsilon}} \times S_r \times \prod_{\lambda} S_{r_{\lambda}}$, where S_n denotes the symmetric group in n letters. Then G acts on $\mathbb{C}[\Sigma]$ by permuting the linear forms $w_{\varepsilon,a}, x_b, y_{\lambda,c}$ in the obvious manner. Consider the following correspondence:

$$\begin{aligned} w_{\varepsilon,a} &\longleftrightarrow T\text{-Chern roots of } Q_{N_{\varepsilon}} \\ x_b &\longleftrightarrow T\text{-Chern roots of } Q_{F_0} \\ y_{\lambda,c} &\longleftrightarrow T\text{-Chern roots of } Q_{F_{\lambda}} \\ d_i &\longleftrightarrow c_1(Q_{\Delta_i})_T \\ e_j &\longleftrightarrow c_1(Q_{\xi_j})_T \end{aligned}$$

Such a correspondence defines a natural map $\rho : \mathbb{C}[\Sigma]^G \rightarrow H_T^*(X^*)$.

Let $\phi : \widetilde{X}^* \rightarrow X^*$ denote the blow-up of X^* along Z with exceptional divisor E . Define $\widetilde{\Sigma}$ to be the polyhedral complex obtained from Σ by adding the ray through the vector $\sum x_b + \sum y_{\lambda,c} + \sum d_i + \sum e_j$. As before, let $w_{\varepsilon,a}, x_b, y_{\lambda,c}, d_i$, and e_j denote the linear forms on $\mathbb{C}[\widetilde{\Sigma}]$ which are dual to vectors $w_{\varepsilon,a}, x_b, y_{\lambda,c}, d_i$, and e_j . Let t be the linear form dual to $\sum x_b + \sum y_{\lambda,c} + \sum d_i + \sum e_j$. Then:

$$\mathbb{C}[\widetilde{\Sigma}] \cong \mathbb{C}[t, w_{\varepsilon,a}, x_b, y_{\lambda,c}, d_i, e_j] / \prod_{b,c,\lambda,i,j} x_b y_{\lambda,c} d_i e_j.$$

G acts on $\mathbb{C}[\widetilde{\Sigma}]$ in the obvious manner. Consider the correspondence:

$$\begin{aligned} w_{\varepsilon,a} &\longleftrightarrow T\text{-Chern roots of } \phi^* Q_{N_{\varepsilon}} \\ x_b &\longleftrightarrow T\text{-Chern roots of } \phi^* Q_{F_0} \otimes \mathcal{O}(-E) \\ y_{\lambda,c} &\longleftrightarrow T\text{-Chern roots of } \phi^* Q_{F_{\lambda}} \otimes \mathcal{O}(-E) \\ d_i &\longleftrightarrow c_1(\phi^* Q_{\Delta_i} \otimes \mathcal{O}(-E))_T \\ e_j &\longleftrightarrow c_1(\phi^* Q_{\xi_j} \otimes \mathcal{O}(-E))_T \end{aligned}$$

By theorem 8, this correspondence induces a well-defined homomorphism $\rho : \mathbb{C}[\widetilde{\Sigma}]^G \rightarrow H_T^*(X^*)$.

It is easy to see that $\nu^* : \mathbb{C}[\Sigma]^G \rightarrow \mathbb{C}[\widetilde{\Sigma}]^G$ and similarly $\nu_* : \mathbb{C}[\widetilde{\Sigma}]^G \rightarrow \mathbb{C}[\Sigma]^G$. We have the following important lemmas:

Lemma 10.

$$(8) \quad \rho\nu^* = \phi^*\rho$$

$$(9) \quad \phi_*\rho = \rho\nu_*$$

Proof. For notational convenience, let $\bar{w}_{\varepsilon,a}, \bar{x}_b, \dots$ denote the T -Chern roots corresponding to $w_{\varepsilon,a}, x_b, \dots$. With this notation, $\rho(f)(w_{\varepsilon,a}, x_b, \dots)$ is equal to $f(\bar{w}_{\varepsilon,a}, \bar{x}_b, \dots)$. We first prove 8: We have:

$$\begin{aligned} \phi^*\rho(f) &= f(\phi^*\bar{w}_{\varepsilon,a}, \phi^*\bar{x}_b, \dots) \\ &= f(\bar{w}_{\varepsilon,a}, \bar{x}_b + \bar{t}, \dots) = f(\rho(w_{\varepsilon,a}), \rho(x_b + t), \dots) \\ &= f(\rho\nu^*w_{\varepsilon,a}, \rho\nu^*x_b, \dots) = \rho\nu^*f \end{aligned}$$

8 allows us to reduce the proof of 9 to the case where $f = t^n$. Let $N = \text{codim } Z$. For $n < N-1$, $\nu_*t^n = 0 = \phi_*\rho(t)^n$, so the claim is true in this case. Otherwise, for $n = (\ell - 1 + j)$, $\phi_*\rho(x_0^n) = (-1)^{j-1}(i_Z)_*s_j(N_{Z/X^*})_T = (-1)^{j-1}(i_Z)_*i_Z^*s_j(M)_T = (-1)^{j-1}s_j(M)_{Tc_{top}(M)_T}$. Here $s_j(M)_T$ denotes the j -th equivariant Segre class of the bundle $M = Q_{F_0} \oplus \bigoplus_{\lambda} Q_{F_{\lambda}} \oplus \bigoplus_{\Delta_i} Q_{\Delta_i} \oplus \bigoplus_j Q_{\xi_j}$. The last equality follows from lemma 7. Clearly this last expression is equal to $\rho(P)$ for some universal polynomial P in the Chern roots of M . We are therefore reduced to proving that $\nu_*(f) = P$. Let $s = \text{rank}(N_{W/Z})$ and let $\mu : \widetilde{\mathbb{C}^{s+N}} \rightarrow \mathbb{C}^{s+N}$ denote the blow-up of $\mathbb{C}^{s+N}$ along $\mathbb{C}^s \times 0$, where we give both spaces the structure of toric varieties, with big torus L . $\mathbb{C}[\widetilde{\Sigma}]$ and $\mathbb{C}[\Sigma]$ both correspond to the rings of piece-wise polynomial functions on the fans of $\widetilde{\mathbb{C}^{s+N}}$ and $\mathbb{C}^{s+N}$, which are in turn isomorphic to the L -equivariant cohomology rings of the above spaces. We may view ν_* as the polyhedral version of the equivariant pushforward $\mu_* : H_L^*(\widetilde{\mathbb{C}^{s+N}}) \rightarrow H_L^*(\mathbb{C}^{s+N})$. Then $\nu_*f = P$ follows from the fact that $\mu_*(c_1(E)_L)^{N-1+j} = P$, where here we evaluate P at the equivariant Chern classes of the coordinate hyperplanes. $\square$

Remark 6. As in section 5.3, the above theorems hold for piece-wise convergent power series.

10. BLOW-UP FORMULA FOR ORBIFOLD ELLIPTIC GENUS

Let X be a smooth projective variety with a holomorphic $G \times T$ action, $D = \sum_I \alpha_i D_i$ a $G \times T$ -invariant G -normal crossing divisor with coefficients $\alpha_i < 1$, $g, h \in G$ a pair of commuting elements, and $X_{\gamma}^{g,h}$ a component of the common fixed point locus of g and h . Recall the definition given in section 3 for the orbifold elliptic class of the pair $(X_{\gamma}^{g,h}, D)$. The goal in this section is to prove the following theorem:

Theorem 8. *Let $f : \tilde{X} \rightarrow X$ be the blow-up along a smooth $G \times T$ -invariant subvariety Z which has normal crossings with respect to the components of D . Fix a commuting pair $g, h \in G$ and a component $X_\gamma^{g,h}$ of the fixed point locus $X^{g,h}$. Let $\{\tilde{X}_\mu^{g,h}\}$ denote the components of $\tilde{X}^{g,h}$ which get mapped to $X_\gamma^{g,h}$ under f . Then:*

$$f_* \sum_{\mu} \mathcal{E}ll_{orb}(\tilde{X}_\mu^{g,h}, \tilde{D}) = \mathcal{E}ll_{orb}(X_\gamma^{g,h}, D)$$

Here $\tilde{D}$ is the divisor satisfying $f^*(K_X + D) = K_{\tilde{X}} + \tilde{D}$.

Proof. By the projection formula, it suffices to assume that every component of D intersects Z with multiplicity 1. Furthermore, by applying deformation to the normal cone (lemma 6) we may assume that X is the normal cone space X^* described in section 9. Using the notation from section 9, we have the following correspondences:

$$\begin{aligned} H &\longleftrightarrow (g, h) \\ Z \cap X_\gamma^{g,h} &\longleftrightarrow W \\ \{D_j\}_{j \in I_\gamma^{g,h}} &\longleftrightarrow \{V_{\xi_j}\} \\ \{D_i\}_{i \notin I_\gamma^{g,h}} &\longleftrightarrow \{V_{\Delta_i}\} \\ X_\gamma^{g,h} &\longleftrightarrow \bigcap_{\varepsilon} V_{N_\varepsilon} \cap \bigcap_{\lambda} V_{F_\lambda} \cap \bigcap_j V_{\xi_j} \\ Z &\longleftrightarrow V_{F_0} \cap \bigcap_{\lambda} V_{F_\lambda} \cap \bigcap_i V_{\Delta_i} \cap \bigcap_j V_{\xi_j} \end{aligned}$$

Given these identifications, $\mathcal{E}ll_{orb}(X_\gamma^{g,h}, D) =$

$$\begin{aligned} &\prod_{T X^*} \frac{\frac{x_i}{2\pi i} \theta(\frac{x_i}{2\pi i} - z) \theta'(0)}{\theta(\frac{x_i}{2\pi i}) \theta(-z)} \prod_{\lambda, Q_{F_\lambda}} \frac{\theta(\frac{x_{\lambda,c}}{2\pi i}) \theta(\frac{x_{\lambda,c}}{2\pi i} + \lambda(g) - \lambda(h)\tau - z)}{\theta(\frac{x_{\lambda,c}}{2\pi i} - z) \theta(\frac{x_{\lambda,c}}{2\pi i} + \lambda(g) - \lambda(h)\tau)} \times \\ &\prod_{\varepsilon, Q_{N_\varepsilon}} \frac{\theta(\frac{x_{\varepsilon,a}}{2\pi i}) \theta(\frac{x_{\varepsilon,a}}{2\pi i} + \varepsilon(g) - \varepsilon(h)\tau - z)}{\theta(\frac{x_{\varepsilon,a}}{2\pi i} - z) \theta(\frac{x_{\varepsilon,a}}{2\pi i} + \varepsilon(g) - \varepsilon(h)\tau)} \times \\ &\prod_j \frac{\theta(\frac{\xi_j}{2\pi i}) \theta(\frac{\xi_j}{2\pi i} + \lambda_j(g) - \lambda_j(h)\tau - (-\alpha_j + 1)z) \theta(-z)}{\theta(\frac{\xi_j}{2\pi i} - z) \theta(\frac{\xi_j}{2\pi i} + \lambda_j(g) - \lambda_j(h)\tau) \theta(-(-\alpha_j + 1)z)} \times \\ &\prod_i \frac{\theta(\frac{\Delta_i}{2\pi i} - (-\alpha_i + 1)z) \theta(-z)}{\theta(\frac{\Delta_i}{2\pi i} - z) \theta(-(-\alpha_i + 1)z)} \end{aligned}$$

Here $x_{\lambda,c}$, $x_{\varepsilon,a}$, ξ_j , and Δ_i denote the equivariant Chern roots of the bundles, Q_{F_λ} , Q_{N_ε} , Q_{ξ_j} , and Q_{Δ_i} , respectively. The above equality follows from lemma 7.

Let $f : \widetilde{X}^* \rightarrow X^*$ be the blow-up along Z with exceptional divisor E . Before proceeding further, it will be convenient to set up some new notation. Consider the collection $I = \{N_\varepsilon, F_0, F_\lambda, \xi_j, \Delta_i, E\}$. Let $\widetilde{Q}_{N_\varepsilon} = f^*Q_{N_\varepsilon}$. For $A \in I$, $A \neq N_\varepsilon, E$, let $\widetilde{Q}_A = f^*Q_A \otimes \mathcal{O}(-E)$. For ease of notation later on, we also define $\widetilde{Q}_E = \mathcal{O}(E)$. For $A \in I - \{E\}$, we let $\widetilde{V}_A$ denote the proper transform of V_A , and we let $\widetilde{V}_E$ simply equal E .

Let $\widetilde{X}_\mu^{*,g,h}$ be a connected component of $\widetilde{X}^{*,g,h}$ which gets mapped to $X_\gamma^{g,h}$ under f . This space is the complete intersection of subvarieties $\widetilde{V}_A$ for A in some indexing set $I_\mu \subset I$. For $A \in I_\mu$, $\widetilde{Q}_A|_{\widetilde{X}_\mu^{*,g,h}} = N_A \otimes \widetilde{\lambda}_A$ for some (g, h) character $\widetilde{\lambda}_{A,\mu}$ and some (g, h) -trivial bundle $N_{A,\mu}$. We extend the definition of $\widetilde{\lambda}_{A,\mu}$ by letting $\widetilde{\lambda}_{A,\mu} = 0$ for $A \notin I_\mu$.

Finally, we define indices β_A as follows: For $A = \xi_j, \Delta_i$, or E , we let β_A be the coefficient of $\widetilde{V}_A$ as a divisor in $\widetilde{D}$. Otherwise, we let $\beta_A = 0$.

Applying corollary 1, lemma 9, and the above definitions, we obtain the following convenient expression for $\mathcal{E}ll_{orb}(\widetilde{X}_\mu^{*,g,h}, \widetilde{D})$:

$$f^* \left\{ \prod_{T \in X^*} \frac{\frac{x_i}{2\pi i} \theta(\frac{x_i}{2\pi i} - z) \theta'(0)}{\theta(\frac{x_i}{2\pi i}) \theta(-z)} \prod_{m, Q_A, A \in I} \frac{\theta(\frac{a_m}{2\pi i}) \theta(-z)}{\frac{a_m}{2\pi i} \theta(\frac{a_m}{2\pi i} - z) \theta'(0)} \right\} \times \\ \prod_{\ell, \widetilde{Q}_A, A \in I} \frac{\frac{\tilde{a}_\ell}{2\pi i} \theta(\frac{\tilde{a}_\ell}{2\pi i} + \widetilde{\lambda}_{A,\mu}(g) - \widetilde{\lambda}_{A,\mu}(h) \tau - (-\beta_A + 1)z) \theta'(0)}{\theta(\frac{\tilde{a}_\ell}{2\pi i} + \widetilde{\lambda}_{A,\mu}(g) - \widetilde{\lambda}_{A,\mu}(h) \tau) \theta(-(-\beta_A + 1)z)}$$

For $A \in I - \{E\}$ we define, for convenience of notation, $\lambda_A \in R((g, h))$ and $\alpha_A \in \mathbb{Q}$ as follows: For $A = N_\varepsilon, F_\lambda, \xi_j$, and Δ_i , we let $\lambda_A = \varepsilon, \lambda, \lambda_j$, and λ_i , respectively. Otherwise, we set $\lambda_A = 0$. Next, for $A = \xi_j$ or Δ_i , we let $\alpha_A = \alpha_j$ or α_i . Otherwise, we set $\alpha_A = 0$.

By the projection formula, we are reduced to proving the following formula:

$$(10) \quad f_* \sum_\mu \prod_{\ell, \widetilde{Q}_A, A \in I} \frac{\frac{\tilde{a}_\ell}{2\pi i} \theta(\frac{\tilde{a}_\ell}{2\pi i} + \widetilde{\lambda}_{A,\mu}(g) - \widetilde{\lambda}_{A,\mu}(h) \tau - (-\beta_A + 1)z) \theta'(0)}{\theta(\frac{\tilde{a}_\ell}{2\pi i} + \widetilde{\lambda}_{A,\mu}(g) - \widetilde{\lambda}_{A,\mu}(h) \tau) \theta(-(-\beta_A + 1)z)} =$$

$$(11) \quad \prod_{m, Q_A, A \in I - E} \frac{\frac{a_m}{2\pi i} \theta(\frac{a_m}{2\pi i} + \lambda_A(g) - \lambda_A(h) \tau - (-\alpha_A + 1)z) \theta'(0)}{\theta(\frac{a_m}{2\pi i} + \lambda_A(g) - \lambda_A(h) \tau) \theta(-(-\alpha_A + 1)z)}$$

Naturally, $\tilde{a}_\ell = \tilde{a}_\ell(A)$ denote the equivariant Chern roots of $\widetilde{Q}_A$ and $a_m = a_m(A)$ denote the equivariant Chern roots of Q_A .

Referring to the notation from section 9 for $A = N_\varepsilon, F_0, F_\lambda, \Delta_k, \xi_j$ and $i = 1, \dots, \text{rank}(A)$, let us define $x_{A,i} \in \mathbb{C}[\Sigma] = w_{\varepsilon,i}, x_i, y_{\lambda,i}, d_k, e_j$, respectively.

For $A \in I$ and $i = 1, \dots, \text{rank}(A)$ we define $\tilde{x}_{A,i} \in \mathbb{C}[\tilde{\Sigma}]$ similarly. Define $F \in \mathbb{C}[[\Sigma]]$ to be the power series:

$$F = \prod_{i,A \in I-E} \frac{\frac{x_{A,i}}{2\pi i} \theta(\frac{x_{A,i}}{2\pi i} + \lambda_A(g) - \lambda_A(h)\tau - (-\alpha_A + 1)z) \theta'(0)}{\theta(\frac{x_{A,i}}{2\pi i} + \lambda_A(g) - \lambda_A(h)\tau) \theta(-(-\alpha_A + 1)z)}$$

Define $F_\mu \in \mathbb{C}[[\tilde{\Sigma}]] =$

$$\prod_{i,A \in I} \frac{\frac{\tilde{x}_{A,i}}{2\pi i} \theta(\frac{\tilde{x}_{A,i}}{2\pi i} + \tilde{\lambda}_{A,\mu}(g) - \tilde{\lambda}_{A,\mu}(h)\tau - (-\beta_A + 1)z) \theta'(0)}{\theta(\frac{\tilde{x}_{A,i}}{2\pi i} + \tilde{\lambda}_{A,\mu}(g) - \tilde{\lambda}_{A,\mu}(h)\tau) \theta(-(-\beta_A + 1)z)}$$

Clearly F is ρ applied to expression 10 and F_μ is ρ applied to the μ -th summand in expression 11. By lemma 10, we are reduced to proving $\nu_* \sum_\mu F_\mu = F$. To do this, think of (Σ, N_Σ) as the polyhedral complex associated to the toric variety $\mathbb{C}^M$ where $M = \dim \Sigma$. Let $\widetilde{\mathbb{C}^M}$ be the toric blow-up of $\mathbb{C}^M$ which corresponds to the polyhedral subdivision $\tilde{\Sigma} \rightarrow \Sigma$ described in section 9. We may view g and h as elements of the big torus of $\mathbb{C}^M$, i.e., as elements of a finite index sup-lattice of N_Σ . Under this identification, $x_{A,i}(g) = \lambda_A(g)$.

The (g, h) -fixed components of $\widetilde{\mathbb{C}^M}$ are in one-one correspondence with the fixed components $\widetilde{X}_\mu^{*,g,h}$ and in one-one correspondence with subcones $C_\mu \subset \tilde{\Sigma}$. These are the cones of maximal dimension which correspond to affine open sets U_{C_μ} of the form $\mathbb{C}^a \times (\mathbb{C}^*)^b$, where the characters of the (g, h) -representation $\mathbb{C}^a$ are all non-trivial. For $C \supset C_\mu$, let $I(C)$ index the collection of piece-wise linear functions $\tilde{x}_{A,i}$ which are dual to C . Since g and $h \in N_\Sigma$ lie inside C_μ , it makes sense to evaluate $\tilde{x}_{A,i}|_C$ at g and h . One sees easily in fact that $\tilde{x}_{A,i}|_C(g) = \tilde{\lambda}_{A,\mu}(g)$ and similarly for $\tilde{x}_{A,i}|_C(h)$. When distinguishing between different cones, it will be convenient to denote the collection $\{\tilde{x}_{A,i}|_C\}_{I(C)}$ by $\{\tilde{x}_{C,j}\}_{j=1}^{|I(C)|}$. With this notation, if $\tilde{x}_{C,j} = \tilde{x}_{A,i}|_C$, we also define $\beta_j = \beta_A$. We define $x_{C,j}$ and α_j similarly when $C \subset \Sigma$.

From this it follows that for $C \supset C_\mu$:

$$F_\mu|_C = \prod_{j=1}^{\dim C} \frac{\frac{\tilde{x}_{C,j}}{2\pi i} \theta(\frac{\tilde{x}_{C,j}}{2\pi i} + \tilde{x}_{C,j}(g) - \tilde{x}_{C,j}(h)\tau - (-\beta_j + 1)z) \theta'(0)}{\theta(\frac{\tilde{x}_{C,j}}{2\pi i} + \tilde{x}_{C,j}(g) - \tilde{x}_{C,j}(h)\tau) \theta(-(-\beta_j + 1)z)}$$

Otherwise, if C_μ is not contained in C , it is easy to see that $F_\mu|_C = 0$.

Now let $C_\gamma \subset \Sigma$ be the cone which corresponds to $(X^*)_\gamma^{g,h}$. Fix a cone $K \subset \Sigma$ containing C_γ . Let $\tilde{\Sigma}_K$ denote the subdivision of K inside $\tilde{\Sigma}$. Each cone $C \subset \tilde{\Sigma}_K$ with the same dimension as K contains a unique cone C_μ for some μ . Moreover, every cone C_μ is contained in one such cone $C \subset \tilde{\Sigma}_K$. We therefore have that

$$\sum_{\mu} F_{\mu}|_C = \prod_{j=1}^{\dim K} \frac{\theta(\frac{\tilde{x}_{C,j}}{2\pi i} + \tilde{x}_{C,j}(g) - \tilde{x}_{C,j}(h)\tau - (-\beta_j + 1)z)\theta'(0)}{\theta(\frac{\tilde{x}_{C,j}}{2\pi i} + \tilde{x}_{C,j}(g) - \tilde{x}_{C,j}(h)\tau)\theta(-(-\beta_j + 1)z)}$$

To complete the proof, it remains to show that $(\nu_* \sum F_{\mu})|_K = F|_K$. By the push-forward formula for ν_* , this is equivalent to proving the identity:

$$\begin{aligned} & \sum_{C \in \tilde{\Sigma}_K} \prod_{j=1}^{\dim K} \frac{\theta(\frac{\tilde{x}_{C,j}}{2\pi i} + \tilde{x}_{C,j}(g) - \tilde{x}_{C,j}(h)\tau - (-\beta_j + 1)z)\theta'(0)}{\theta(\frac{\tilde{x}_{C,j}}{2\pi i} + \tilde{x}_{C,j}(g) - \tilde{x}_{C,j}(h)\tau)\theta(-(-\beta_j + 1)z)} = \\ & \prod_{j=1}^{\dim K} \frac{\theta(\frac{x_{K,j}}{2\pi i} + x_{K,j}(g) - x_{K,j}(h)\tau - (-\alpha_j + 1)z)\theta'(0)}{\theta(\frac{x_{K,j}}{2\pi i} + x_{K,j}(g) - x_{K,j}(h)\tau)\theta(-(-\alpha_j + 1)z)} \end{aligned}$$

Here the functions $\tilde{x}_{C,j}$ are regarded as linear combinations of the functions $x_{K,j}$. The above formula follows from theorem 7 of the preprint [17] or by lemma 8.1 of [7]. This completes the proof. $\square$

Now, let Z be a projective $\mathbb{Q}$ -Gorenstein variety with log terminal singularities, and a regular $G \times T$ action. Let $f : X \rightarrow Z$ and $g : Y \rightarrow Z$ be two equivariant resolutions of singularities. We assume that the exceptional locus of both resolutions is a G -normal divisor with simple normal crossings. Define D_X so that $K_X + D_X = f^*K_Z$; define D_Y similarly. Then the equivariant orbifold elliptic genera of (X, D_X) and (Y, D_Y) coincide. Indeed, by the equivariant version of the weak factorization theorem [1], we may connect X to Y by a sequence of equivariant blow-ups and blow-downs in such a way that the blow-ups at each intermediate pair (X_i, D_{X_i}) occur along a smooth base with normal crossings with respect to the components of D_{X_i} . Moreover, the procedure described in [7] theorem 3.7 to make the intermediate pairs (X_i, D_{X_i}) G -normal extends to the T -equivariant case. Hence, by the equivariant change of variables formula, $Ell_{orb}(X, D_X, G) = Ell_{orb}(Y, D_Y, G)$.

We will, however, require a slightly stronger version of the above result for the purposes of this paper:

Theorem 9. *Let $f : (X, D_X) \rightarrow (Y, D_Y)$ be a $G \times T$ -equivariant birational morphism between smooth, equivariant, G -normal log terminal pairs (X, D_X) and (Y, D_Y) . Assume furthermore that $f^*(K_Y + D_Y) = K_X + D_X$. Then $f_* Ell_{orb}(X, D_X, G) = Ell_{orb}(Y, D_Y, G)$.*

Proof. The weak factorization theorem allows us to factor f into a sequence of equivariant blow-ups and blow-downs

$$X = X_0 \dashrightarrow X_1 \dashrightarrow \cdots \dashrightarrow X_k = Y$$

such that for some intermediate index i_0 the maps $X_i \rightarrow X$ are morphisms for $i \leq i_0$ and the maps $X_i \rightarrow Y$ are morphisms for $i \geq i_0$. Moreover, by [7] theorem 3.7, we may still guarantee that all the intermediate varieties are G -normal with respect to the appropriate divisors. Note that since f is itself a smooth morphism, we may conclude that the maps $X_i \rightarrow Y$ are smooth morphisms for all i .

We now apply induction on k . For $k = 1$, the theorem is obvious. Otherwise, consider the intermediate variety X_1 . By assumption, $X_1 \neq X, Y$. Either $X \dashrightarrow X_1$ is a blow-up or blow-down. Suppose first that $X \leftarrow X_1$ is a blow-down. Call this morphism g . Define D_1 so that $K_{X_1} + D_1 = g^*(K_X + D_X)$. Note that if $h : X_1 \rightarrow Y$ is the morphism $f \circ g$, then $K_{X_1} + D_1 = h^*(K_Y + D_Y)$. By the change of variables formula, $\mathcal{E}ll_{orb}(X, D_X, G) = g_* \mathcal{E}ll_{orb}(X_1, D_1, G)$. Therefore $f_* \mathcal{E}ll_{orb}(X, D_X, G) = h_* \mathcal{E}ll_{orb}(X_1, D_1, G)$. But this is equal to $\mathcal{E}ll_{orb}(Y, D_Y, G)$ by the induction hypothesis.

The case in which $X \rightarrow X_1$ is a blow-down is proved similarly. $\square$

11. THE EQUIVARIANT MCKAY CORRESPONDENCE

Here we use the results from the preceding sections to prove an equivariant analogue of the McKay Correspondence for the elliptic genus. As a corollary, we will arrive at an equivariant version of the DMVV formula.

Let X be a projective variety with a $G \times T$ action, where G is a finite group and T is a compact torus. Let $\psi : X \rightarrow X/G$ be the quotient morphism. Assume that X/G has an equivariant crepant resolution V and that $\psi^*(K_{X/G}) = K_X$. Let $F \subset X$ be a fixed component of the $G \times T$ -action, and let $\{P\} \subset V$ denote the fixed components in V which get mapped to $\psi(F)$. Then:

Theorem 10.

$$\int_F \frac{\mathcal{E}ll_{orb}(X, G)}{e(F)} = \sum_P \int_P \frac{\mathcal{E}ll(V)}{e(P)}.$$

Proof. Let $Z \rightarrow V$ be a sequence of equivariant blow-ups of V so that the exceptional locus of the resolution $\pi : Z \rightarrow X/G$ is a divisor with simple normal crossings. Let D_Z be the divisor on Z such that $K_Z + D_Z = \pi^* K_{X/G}$. Define $\hat{Z}_0$ to be the normalization of Z in the function field of X . By Abhyankar's Lemma, the induced map $\mu_0 : \hat{Z}_0 \rightarrow Z$ is a toroidal morphism of toroidal embeddings. Let $\hat{Z}$ be a projective toroidal resolution of singularities of $\hat{Z}_0$, and define $D_{\hat{Z}}$ so that $K_{\hat{Z}} + D_{\hat{Z}} = \mu^*(K_Z + D_Z)$, where $\mu : \hat{Z} \rightarrow Z$ is the obvious map. We may further assume that $\hat{Z}$ has a

G action, and that the pair $(\hat{Z}, D_{\hat{Z}})$ is G -normal. (see [2]). We obtain the following commutative diagram:

$$\begin{array}{ccc} \hat{Z} & \xrightarrow{\mu} & Z \\ \phi \downarrow & & \downarrow \pi \\ X & \xrightarrow{\psi} & X/G \end{array}$$

Here, the vertical arrows are resolutions of singularities, and the horizontal arrows are birational to a quotient by G . It is clear that all the constructions involved (normalization, blow-up along a T -invariant ideal sheaf) are T -equivariant, and that consequently the above morphisms are T -equivariant.

Since $\phi^*(K_X) = K_{\hat{Z}} + D_{\hat{Z}}$, we have that $\mathcal{E}ll_{orb}^T(X, G) = \phi_* \mathcal{E}ll_{orb}(\hat{Z}, D_{\hat{Z}}, G)$. Let $\{L\} \subset \hat{Z}$ denote the fixed components of $\hat{Z}$ which map to F . By functorial localization:

$$\int_F \frac{\mathcal{E}ll_{orb}(X, G)}{e(F)} = \sum_L \int_L \frac{\mathcal{E}ll_{orb}(\hat{Z}, D_{\hat{Z}}, G)}{e(L)}.$$

Now, by theorem 6, $\mu_* \mathcal{E}ll_{orb}(\hat{Z}, D_{\hat{Z}}, G) = \mathcal{E}ll(Z, D_Z)$. Let $\{K\} \subset Z$ denote the fixed components which get mapped to $\psi(F)$ under the resolution $Z \rightarrow X/G$. Clearly $\{L\} = \phi^{-1}F = \phi^{-1}\psi^{-1}\psi(F) = \mu^{-1}\pi^{-1}\psi(F) = \mu^{-1}\{K\}$. Thus, by functorial localization applied to μ_* , we have:

$$\sum_L \int_L \frac{\mathcal{E}ll_{orb}(\hat{Z}, D_{\hat{Z}}, G)}{e(L)} = \sum_K \int_K \frac{\mathcal{E}ll(Z, D_Z)}{e(K)}.$$

Finally, since $\{K\}$ denotes the fixed components of Z which get mapped to $\{P\}$, functorial localization applied to $Z \rightarrow V$ completes the proof. $\square$

We now discuss some corollaries of the above result. We begin with a proof of the equivariant DMVV conjecture. Let $\mathbb{P}_2^{(n)} = (\mathbb{P}_2)^n / S_n$ denote the n th symmetric product of the projective plane. The natural group action of $T = S^1 \times S^1$ on $\mathbb{P}_2$ extends to $(\mathbb{P}_2)^n$ in the obvious manner, and commutes with the action of S_n . The action of T on $\mathbb{P}_2$ also extends to $\mathbb{P}_2^{[n]}$, the Hilbert scheme of n points on $\mathbb{P}_2$, and the Hilbert-Chow morphism $\mathbb{P}_2^{[n]} \rightarrow \mathbb{P}_2^{(n)}$ is an equivariant crepant resolution.

Sitting inside $\mathbb{P}_2^{(n)}$ is the open variety $(\mathbb{C}^2)^{(n)}$. It has a single T -fixed point p , which is the image under the quotient morphism of the $S_n \times T$ fixed point $(0, 0) \times \dots \times (0, 0) \in (\mathbb{C}^2)^n$. The pre-image of $(\mathbb{C}^2)^{(n)}$ under the Hilbert-Chow morphism is just $(\mathbb{C}^2)^{[n]}$. Hence, the pre-image of p under the Hilbert-Chow morphism is simply the collection of T -fixed points of $(\mathbb{C}^2)^{[n]}$. If (u_1, u_2)

denote the equivariant parameters of the T -action, let $t_j = e^{2\pi i u_j}$. Then the above theorem implies that:

$$Ell_{orb}((\mathbb{C}^2)^n, S_n, t_1, t_2) = Ell((\mathbb{C}^2)^{[n]}, t_1, t_2).$$

Note that the LHS involves equivariant data associated to the single fixed point p , whereas the RHS involves a sum of equivariant data associated to partitions of n .

For z the complex parameter appearing in the definition of the elliptic class and τ the lattice parameter used in the definition of the Jacobi theta function, let $y = e^{2\pi i z}$ and $q = e^{2\pi i \tau}$. Viewing the equivariant elliptic indices as formal power series in the variables q, y, t_1 , and t_2 , and applying theorem 3.1 of [14], we have the following equivariant analogue of the DMVV formula:

Theorem 11. *Write $Ell(\mathbb{C}^2, t_1, t_2) = \sum_{m \geq 0, \ell, k} c(m, \ell, k) q^m y^\ell t_1^{k_1} t_2^{k_2}$. Then*

$$\sum_{n > 0} p^n Ell((\mathbb{C}^2)^{[n]}, t_1, t_2) = \prod_{m \geq 0, n > 0, \ell, k} \frac{1}{(1 - p^n q^m y^\ell t_1^{k_1} t_2^{k_2})^{c(nm, \ell, k)}}$$

We next discuss the equivariant elliptic genus analogue of the classical McKay correspondence, which was originally proved in [17]. Let $G \subset SU(2)$ be a finite subgroup. G acts on $\mathbb{C}^2$ in the obvious fashion, and commutes with the diagonal action of $T = S^1$. Let $V \rightarrow \mathbb{C}^2/G$ be the crepant resolution of singularities. The action of T lifts to V , and the fixed components of this action are compact.

Theorem 12. $Ell_{orb}(\mathbb{C}^2, G, t) = Ell(V, t)$

Proof. View $\mathbb{C}^2$ as an open subset of $\mathbb{P}^2$, with the action of G and T extending to $\mathbb{P}^2$ in the obvious manner. The space $\mathbb{P}^2/G$ still has only an isolated singularity at the image of the origin $[0 : 0 : 1]$. Hence $\mathbb{P}^2/G$ has an equivariant crepant resolution which is a compactification of V . The above theorem then follows by letting $F = (0, 0)$ in theorem 10. $\square$

12. APPENDIX

Lemma 11. *Let $f : X \rightarrow Y$ be a T -map of smooth compact simply connected Kahler varieties. Let $D \subset Y$ be a T -invariant divisor and let E_i be T -invariant normal crossing divisors on X such that $f^*D = \sum a_i E_i$ as Cartier divisors. Then for any ε -regular neighborhood U_ε of D there exist generators $\Theta_{E_i}^T$ for $c_1^T(E_i)$ and Θ_D^T for $c_1^T(D)$ with the following properties:*

- (1) Θ_D^T has compact support in U_ε and $\Theta_{E_i}^T$ have compact support in $f^{-1}(U_\varepsilon)$.

(2) $f^*\Theta_D^T = \sum a_i \Theta_{E_i}^T + d_T(\eta)$ on the level of forms, where η is a T -invariant form with compact support in U_ε .

(3) Θ_D^T and $\Theta_{E_i}^T$ represent the equivariant Thom classes of the varieties D and E_i

The only real issue above is to ensure that η has compact support in the desired neighborhood.

Proof. We first solve this problem in the non-equivariant category. For V any Cartier divisor, denote by L_V the line bundle it induces. Let U_ε be a T -invariant tubular neighborhood of D of radius ε . Outside U_ε , the constant function 1 is a section of L_D . Define a metric h_{far} in this region by $h_{far} = ||1||^2 \equiv 1$. Let h_{near} be a metric inside U_ε . Piece the two metrics into a global metric h on L_D using a partition of unity. The first Chern class of L_D is then represented by the form $\Theta_D = \frac{i}{2\pi} \bar{\partial} \partial \log h$. This form clearly has compact support in U_ε .

Let U_{ε_i} be tubular neighborhoods of E_i . Choose ε_i small enough so that each of these neighborhoods is contained in $f^{-1}U_\varepsilon$. Define metrics h_i on E_i in a manner analogous to the above construction of h . Clearly the forms $\Theta_{E_i} = \frac{i}{2\pi} \bar{\partial} \partial \log h_i$ have compact support in U_{ε_i} and represent the first Chern classes of L_{E_i} .

We have two natural choices for a metric on f^*L_D , namely f^*h and $h_1^{a_1} \cdots h_k^{a_k}$. Choose a smooth nonzero function φ so that $f^*h = \varphi h_1^{a_1} \cdots h_k^{a_k}$. Notice that $\varphi \equiv 1$ outside $f^{-1}U_\varepsilon$. We have:

$$\bar{\partial} \partial \log f^*h = \bar{\partial} \partial \log \varphi + \sum_i a_i \bar{\partial} \partial \log h_i.$$

But this implies that $f^*\Theta_D = \sum_i a_i \Theta_{E_i} + \frac{i}{2\pi} \bar{\partial} \partial \log \varphi$. If we let $d^c = \frac{i}{4\pi} (\bar{\partial} - \partial)$, we may write this last equation as:

$$f^*\Theta_D = \sum_i a_i \Theta_{E_i} - dd^c \log \varphi.$$

The form $\eta = -d^c \log \varphi$ clearly has compact support in $f^{-1}U_\varepsilon$. It remains to argue that Θ_D and Θ_{E_i} represent the Thom classes of D and E_i . Suppose V is any effective Cartier divisor on a smooth simply connected compact Kahler variety. Then L_V is nontrivial as a holomorphic line bundle and therefore $c_1(L_V) \neq 0$. This follows from the fact that $H^1(X, \mathcal{O}) = 0$ and therefore that the map $c_1 : H^1(X, \mathcal{O}^*) \rightarrow H^2(X, \mathbb{Z})$ is injective. Furthermore, $c_1(L_V)$ is clearly a non-torsion class in $H^2(X, \mathbb{Z})$. Hence we can find an ω such that $\int_X c_1(L_V) \wedge \omega \neq 0$. Since $c_1(L_V)$ is Poincaré dual to V , we have $\int_X c_1(L_V) \wedge \omega = \int_V \omega = \int_X \Phi_V \wedge \omega$ where Φ_V is the Thom class of the normal

bundle N_V to V . However, if Θ_V is a representative of $c_1(L_V)$ with compact support in N_V , we must have $\Theta_V = a\Phi_V + d\psi$ for some form ψ with compact support in N_V . Therefore $\int_X \Theta_V \wedge \omega = a \int_X \Phi_V \wedge \omega = \int_X \Phi_V \wedge \omega \neq 0$. It follows that $a = 1$. Thus, Θ_D and Θ_{E_i} indeed represent the Thom classes of D and E_i . I should remark that it is well-known that Θ_V is Poincaré dual to V . However, under weaker conditions, a form that is Poincaré dual to a submanifold V may not necessarily represent the Thom class of N_V . For example, if V were homologous to zero, then 0 would certainly be Poincaré dual to V but would not represent its Thom class, which would be non-trivial.

This completes the non-equivariant portion of the proof.

By averaging over the group T , we may assume that all the forms above are T -invariant. For notational simplicity, let us assume that $T = S^1$. Let V be the vectorfield on X induced by the T -action. Let g_i be the functions compactly supported in $f^{-1}U_\varepsilon$ which satisfy the moment map equation $i_V \Theta_{E_i} = dg_i$. Similarly, let W be the vectorfield on Y defined by the T -action and define g so that it satisfies $i_V \Theta_D = dg$ and has support inside U_ε . Note that since f is T -equivariant, $i_V f^* \Theta_D = f^* i_W \Theta_D = f^* dg$. We then have $d(g \circ f) = \sum_i a_i dg_i + i_V d\eta = \sum_i a_i dg_i - di_V \eta$. Hence $g \circ f = \sum_i a_i g_i - i_V \eta$. But this implies that:

$$f^*(\Theta_D + g) = \sum_i a_i (\Theta_{E_i} + g_i) + (d - i_V)\eta.$$

But this is precisely the relation we wish to in the equivariant cohomology. $\square$

Lemma 12. (Excess Intersection Formula) *Let X be a smooth compact variety with irreducible normal crossing divisors $D_1, \dots, D_k$. For $I \subset \{1, \dots, k\}$ denote by $X_{I,j}$ the j th connected component of $\cap_I D_i$ and by $\Phi_{I,j}$ its Thom class. Fix irreducible subvarieties X_{I_1,j_1} and X_{I_2,j_2} . For $I_0 = I_1 \cup I_2$, let $X_{I_0,j}$ be the irreducible components of $X_{I_1,j_1} \cap X_{I_2,j_2}$. Then:*

$$\Phi_{I_1,j_1} \wedge \Phi_{I_2,j_2} = \sum_{I_0,j} \Phi_{I_0,j} \prod_{I_1 \cap I_2} \Phi_i.$$

Proof. Let $N_{I,j}$ be tubular neighborhoods of $X_{I,j}$ which are disjoint for each indexing set I and which satisfy $N_{I,j} \subset N_{I',j'}$ for $X_{I,j} \subset X_{I',j'}$. If we choose Φ_i to have compact support in a sufficiently small tubular neighborhood of D_i , then $\prod_I \Phi_i$ will have compact support in $\prod_j N_{I,j}$. Moreover, the extension by zero of $(\prod_I \Phi_i)|_{N_{I,j}}$ will represent the Thom class of $X_{I,j}$ (see [BT]). We may also ensure that $\Phi_{I_1,j_1} \wedge \Phi_{I_2,j_2}$ has compact support in $\prod_j N_{I_0,j}$. Thus:

$$\begin{aligned}\Phi_{I_1, j_1} \wedge \Phi_{I_2, j_2} &= \sum_{I_0, j} \left(\prod_{I_1} \Phi_i \prod_{I_2} \Phi_i \right) |_{N_{I_0, j}} = \sum_{I_0, j} \left(\prod_{I_0} \Phi_i \prod_{I_1 \cap I_2} \Phi_i \right) |_{N_{I_0, j}} = \\ &= \sum_{I_0, j} \left(\prod_{I_0} \Phi_i \right) |_{N_{I_0, j}} \prod_{I_1 \cap I_2} \Phi_i.\end{aligned}$$

This yields the desired formula. $\square$

Remark 7. Note that the above proof clearly extends to the equivariant category.

REFERENCES

- [1] D. Abramovich, K. Karu, K. Matsuki, J. Włodarczyk, *Torification and factorization of birational maps*, J. Amer. Math. Soc. 15, no. 3, pp. 531-572, 2002.
- [2] D. Abramovich, J. Wang, *Equivariant resolution of singularities in characteristic 0* Math. Res. Lett. 4, no. 2-3, pp. 427-433, 1997.
- [3] M.F. Atiyah, R. Bott, *The moment map and equivariant cohomology*, Topology Vol. 23, No. 1, pp. 1-28, 1984.
- [4] M. Atiyah, I. Singer, *The index of elliptic operators III*, Ann. of Math., (2) 87, pp. 546-604, 1968.
- [5] V.V. Batyrev, *Non-Archimedean integrals and stringy Euler numbers of log-terminal pairs*, J. Eur. Math. Soc. (JEMS), 1, pp. 5-33, 1999.
- [6] L. Borisov, A. Libgober, *Elliptic genera of singular varieties*, Duke Math. J. 116, no. 2, pp. 319-351, 2003.
- [7] L. Borisov, A. Libgober, *McKay correspondence for elliptic genera*, Ann. of Math 161, no. 3, pp. 1521-1569, 2005.
- [8] R. Dijkgraaf, G. Moore, E. Verlinde, H. Verlinde, *Elliptic genera of symmetric products and second quantized strings*, Comm. Math. Phys., 185:1, pp. 197-209, 1997.
- [9] W. Fulton, *Intersection theory, second edition*, Springer Verlag, Berlin 1998.
- [10] W. Fulton, *Introduction to toric varieties*, Ann. of Math Studies 131, Princeton University Press, Princeton, N.J., 1993.
- [11] F. Hirzebruch, T. Berger, R. Jung, *Manifolds and modular forms*, with appendices by Nils-Peter Skoruppa and Paul Baum, Aspects of Mathematics, E20. Friedr. Vieweg & Sohn, Braunschweig, 1992.
- [12] B. Lian, K. Liu, S.T. Yau, *Mirror principle I*, Surveys in differential geometry: differential geometry inspired by string theory, 5, pp. 405-454, Int. Press, Boston, MA, 1999.
- [13] B. Lian, K. Liu, S.T. Yau, *Mirror principle II*, Surveys in differential geometry: differential geometry inspired by string theory, 5, pp. 455-509, Int. Press, Boston, MA, 1999.
- [14] J. Li, K. Liu, J. Zhou, *Topological string partition functions as equivariant indices*, Asian J. Math. Vol. 10, No. 1, pp. 081-114, 2006.
- [15] K. Liu, *On elliptic genera and theta-functions*, Topology Vol. 35, no. 3, pp. 617-640, 1996.
- [16] Wang, Chin-Lung, *K-equivalence in birational geometry and characterizations of complex elliptic genera*, J. Algebraic Geom. 12, no. 2, pp. 285-306, 2003.
- [17] R. Waelde, *Equivariant elliptic genera*, preprint, math.AG/0603521.
- [18] E. Witten, *Supersymmetry and Morse theory*, J. Differential Geom. 17 no. 4, pp. 661-692, 1982.

E-mail address: `rwaelde@math.ucla.edu`